THE LIBRARY OF CONGRESS
1800

Class RS125
Book .P176
Copyright No.

COPYRIGHT DEPOSIT.

FAVORITE PRESCRIPTIONS

OF DISTINGUISHED PRACTITIONERS

WITH

NOTES ON TREATMENT

COMPILED FROM THE
PUBLISHED WRITINGS OR UNPUBLISHED RECORDS OF

DRS. FORDYCE BARKER, ROBERTS BARTHOLOW, SAMUEL D. GROSS, AUSTIN FLINT, ALONZO CLARK, ALFRED L. LOOMIS, F. J. BUMSTEAD, T. G. THOMAS, H. C. WOOD, WM. GOODELL, WM. PEPPER, A. JACOBI, J. M. FOTHERGILL, N. S. DAVIS, J MARION SIMS, WM. H. BYFORD, L. A. DUHRING, D. HAYES AGNEW, E. O. JANEWAY, J. M. DA COSTA, J. SOLIS COHEN, GERMAIN SÉE, MEREDITH CLYMER, J. LEWIS SMITH, FLOYD M. CRANDALL, W. H. THOMSON, L. DUNCAN BULKLEY, C. E. BROWN-SEQUARD, M. A. PALLEN, ALEX. J. C. SKENE, GEO. H. FOX, W. A. HAMMOND, E. C. SPITZKA. L. EMMETT HOLT, H. A. HARE, ETC., ETC.

EDITED BY

B. W. PALMER, A.M., M.D.

SEVENTH EDITION

NEW YORK:
E. B. TREAT & COMPANY
241-243 WEST 23D STREET
1901

R S125
P 176

THE LIBRARY OF
CONGRESS,
TWO COPIES RECEIVED
JUN. 6 1901
COPYRIGHT ENTRY
June 6, 1901
CLASS a XXc. No.
10784
COPY B.

COPYRIGHT, 1887–1901.
BY E. B. TREAT & CO.
NEW YORK.

CONTENTS.

I.

PAGE

Diseases of the Nose, Throat, Mouth and Chest..... 7

II.

Diseases of the Abdominal Viscera..... 43

III.

Diseases of the Nervous System..... 71

IV.

Diseases of Women..... 107

V.

Diseases of Children..... 123

VI.

Fevers, Rheumatism, Gout and Malaria..... 149

VII.

Venereal Diseases..... 165

VIII.

PAGE

Tonic Preparations 173

IX.

External Applications 185

X.

Cod Liver Oil Preparations 207

XI.

Miscellaneous Prescriptions 211

Index 239

PREFACE TO THE SEVENTH EDITION.

THE popularity and usefulness of this book of "favorite prescriptions" is amply attested by the fact that six large editions have been printed. This edition (7th) has been revised and greatly enhanced by the addition of many pages of new prescriptions of undoubted value.

It is readily apparent why a book of this kind proves of the greatest help to the busy practitioner, who, surrounded by the voluminous medical publications of the times, is often at a loss to select from the mass of material at his disposal that combination of therapeutical agencies best adapted to meet the indications for treatment in well known conditions of disease.

The editor has striven to place before the profession for constant reference, in a condensed, readily accessible way, matter that could only be gleaned from careful study and association with the medical writers and teachers of the times.

No pretence of originality is made. The book is essentially a compilation. The design being to render more available for every-day use and guidance the practical treasures of medical wisdom which have been gathered from the earnest labors and careful observations of the most distinguished practitioners of the age.

DISEASES OF THE NOSE, THROAT, MOUTH AND CHEST.

Pneumonia.

℞ Ammoniæ muriatis..... ʒ iij.
Antim. et pot. tartrat.....gr. ij.
Morphiæ sulphat.....gr. iij.
Syrupi glycyrrhizæ..... ℥ iv.
M. Sig.—A teaspoonful every two hours.

Dr. N. S Davis.

℞ Hydrarg. chlor. mitis.....
Ipecac. pulveris.....aa gr. vj.
Opii pulveris.....gr. iij.
Sacchar. alb.....gr. xxx.
M. Ft. chart. vj.

Sig.—One powder every four hours alternately with the preceding prescription. At the same time cover the chest with emollient poultices.

Dr. N. S. Davis.

Where a typhoid condition of the system is associated with the pneumonia, bleeding is not well borne. In these cases Dr. Davis has obtained admirable results by the use of the preceding prescriptions.

At the end of twenty-four hours he omits the powders, and if the bowels have not been moved he gives a mild laxative. If the symptoms are not favorably

modified in 3 or 4 days, a blister is placed on the side of the chest most affected.

Should the pulse become soft and frequent, the breathing abdominal and the lips of a leaden hue,

℞ Quiniæ sulph.....gr. ij.
Ammon. carbonat.....gr. iv.

M. Sig.—Take at a dose. If delirium becomes troublesome add 10 minims of chloroform to the ammonia mixture. If there is indication of malarial influence, quinine may be given during the remissions.

Sub-Acute Pleurisy.

℞ Potas. acet.....
Inf. digital.....aa ʒ ij-iv.

Sig.—This amount each day; or

℞ Pulv. digital.....
Pulv. scillæ mar.....
Hydrarg. chlo. mit.....aa gr. x.

M. Et. ft. pil. No. x.

Sig.—One pill thrice daily.

Dr. Alonzo Clark.

The indications for treatment are to subdue the inflammation and promote absorption of the effused fluid. Dr. Clark, to accomplish the first of these effects, uses blisters, three being usually sufficient, selecting three spots and applying only one blister at a time, the second and third not being placed in position until the spot of former application has healed. As a diuretic he uses potassii iodidi xxx grs. a day; if this fail to diminish the fluid, he has resource to the above formulæ. If constitutional effects of mercury

declare themselves he returns again to the potassii iodid. Dr. Clark uses other means, as purgatives, vapor baths, and mild counter-irritants when he thinks them indicated, but, having exhausted all medical means without effect (as is sometimes the case), he either resorts to the trocar or does nothing.

Asthmatic Fits of Emphysema.

℞ Tinct. opii..... ʒ j.
Ætheris..... ʒ ij.

Sig.—Sixty drops every twenty minutes. To each dose may be added:

Tinct. lobeliæ ætheræ.....gtt. xx.

Meridith Clymer, M.D.

This formula has often yielded relief in these sudden attacks of asthma when all other means have failed.

Diuretic in Cardiac Dropsy.

℞ Infus. digital..... ℥ iijss.
Acet. scillæ..... ℥ ss.

M. Sig.—A tablespoonful two or three times a day.

Bartholow.

Diphtheritic Croup.

℞ Potassii chloratis..... ʒ ij.
Syrupi limonis..... ℥ j.
Aquæ..... ℥ iij.

Dose according to age of child. If under two years, a teaspoonful; from two to ten, two teaspoonfuls;

over ten, a tablespoonful, given every three hours or every half hour in urgent cases.

Recommended as almost specific in diphtheritic croup by *Dr T. M. Drysdale.*

In Asthmatic Paroxysm.

℞ Tinct. lobeliæ..... ℥ j.
Ammon. iodidi..... ..... ʒ ij.
Ammon. bromidi..... ..... ʒ iij.
Syr. tolutan..... ℥ iij.

M. Sig.—A teaspoonful every one, two, three or four hours. *Dr. Roberts Bartholow.*

Of this prescription Dr. Bartholow says: "It gives relief in a few minutes, and sometimes the relief is permanent.

Emphysema.

℞ Potass. chlorat..... ..... ʒ jss.
Tinct. belladonnæ..... ℥ jss.
Ext. pruni virg. fluid..... .....
Tinct. cinchonæ co.....aa ℥ ij.

M. Sig.—A dessertspoonful four times a day.

Dr. Da Costa.

Where the emphysema is accompanied by chronic bronchitis and loss of appetite, Dr. Da Costa has found the above prescription the most efficient. He advises also that dry cups be applied to the chest night and morning.

In Bronchial Catarrh.

℞ Acet. scillæ..... ℥ ss.
Ext. ipecac. fl..... ʒ ss.

Tinct. opii deodorat..... ʒ j.
Syr. tolutan..... ʒ x.
M. Sig.—A teaspoonful every two, three or four hours. *Bartholow.*

In Spasmodic Asthma.

℞ Potass. bromid..... ℥ j.
Potass. iodid..... ℥ ss.
Aquæ..... ℥ iv.
M. Sig.—A teaspoonful in sufficient water every half hour or hour. *Dr Bartholow.*

In Chronic Phthisis.

℞ Syr. hypophos. comp..... ℥ iijss.
Acid phosphor. dil..... ℥ ss.
M. Sig.—Teaspoonful three times daily.
Dr. Roberts Bartholow.

The above combination is held by Dr. Bartholow to be very effective therapeutically; it may be advantageously given with cod liver oil, after meals, in chronic phthisis, or the following may be given when the effects of arsenic also are desired:

℞ Syr. calcii lacto-phos..... ℥ iv.
Liq. potass. arsenitis..... ʒ j.
M. Sig.—A dessertspoonful ter die.
Dr. Bartholow.

Acute Catarrh.

℞ Tinct. iodinii..... ℥ ss.
Acid carbolic..... ʒ j.
M. Sig.—Place a small, wide-mouthed bottle, con-

taining a moistened sponge, in a vessel of hot water; drop five to ten drops of the solution on the sponge and as the iodine vapor ascends with the vapor of the water, inhale it. *Bartholow.*

In Summer Catarrh or Hay Asthma.

℞ Potass. iodid..... ℥ j.
Liq. Potass. arsenit..... ʒ j.
Aquæ..... ℥ iv.
M. Sig.—A teaspoonful every four or six hours.

This, given internally, combined with the local use by means of a post nasal syringe of the following formula, will often effect a cure in this obstinate malady:

℞ Tinct. iodinii..... ʒ j.
Acid. carbol.....gtts. x.
Aquæ destillat..... ℥ iv.
M. Sig.—Apply with a post nasal syringe.
Bartholow.

In Acute Coryza and Hay Asthma.

℞ Brominii..... ʒ ss.
Alcoholis..... ℥ iv.
M. Sig.—For inhalation. *Bartholow.*

A small quantity may be placed in a wide-mouthed vial and vaporized by the warmth of the hand. The vapor should be snuffed into the nose.

In Chronic Throat Affections.

℞ Acid. tannici..... ʒ ii.
Spts. vini rect..... ʒ i.
Mist. camph.....ad ℥ x.
M. Sig.—As an astringent gargle. *Bartholow.*

In these throat affections the above as a gargle, combined with a spray of a solution of tannin, ʒ grs. to the ℥ is of great benefit.

Chronic Bronchitis, Emphysema, Spasmodic Asthma, Hay Asthma, and Phthisis.

℞ Potass. arsenit.....gr. xv.
Aquæ destillat..... ℥ j.

Take unsized white paper, thoroughly moisten it with this solution, dry and cut into twenty equal parts and roll each part into a cigarette. Two or three of these are to be smoked daily, inhaling the smoke. *Trousseau.*

The above is also found very beneficial in acute and chronic coryza.

In Chronic Bronchitis.

℞ Ext. eucalypt..... ℥ j.
Ammon. muriat..... ʒ ij.
Ext. glycyrrhizæ..... ℥ ij.
Syr. tolu..... ℥ iij.

M. Sig.—A tablespoonful four or six times a day.
Bartholow.

This forms an excellent expectorant in cases of chronic bronchitis and bronchorrhœa.

Chronic Bronchitis.

℞ Ammonii chloridi..... ʒ ij.
Mist. glycyrrhizæ co..... ..... ℥ iij.

M. Sig.—A dessertspoonful three times a day.
Dr. Da Costa.

This formula is adapted for those cases of bronchitis accompanied by chronic catarrh in old persons.

Acute Bronchitis.

℞ Vini ipecacuanhæ..... ʒ ij.
Liq. potassii citratis..... ℥ iv.
Tinct. opii camphoratæ.....
Syrupi acaciæ.....aa ℥ j.

M. Sig.—A tablespoonful thrice daily in the first stage of ordinary acute bronchitis. *Dr. Da Costa.*

This union of the sedative effects of opium with the excito-secretory action of the ipecacuanha on the congested mucous membrane has been found very serviceable.

Aphthæ of Phthisis.

℞ Quiniæ sulph.....gr. j.
Olei piperis nigris.....gtt. j.
Aquæ..... ℥ j.

M. Sig.—May be applied with a brush, or mouth rinsed with it. This is the wash in use at Roosevelt Hospital, and is at the same time antiseptic and stimulant.

In Mercurial Salivation.

℞ Acidi tannici..... ʒ i.
Mel. rosæ..... ℥ ij.
Aquæ..... ℥ vj.

M. Sig.—As a gargle. *Bartholow.*

Whooping Cough Mixture.

Infants' Hospital, Randall's Island, N. Y.

℞ Acid. nitric. dil.....fl. ʒ i.
Syr. pruni virg.....fl. ℥ ½.
Aquæ q. s. ad.....fl. ℥ 2.
Mix. Dose: a teaspoonful.

Whooping Cough Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Tinct. nucis vom.....fl. ʒ 2.
Vin. ipecac.....fl. ʒ 2½.
Syr. sarsap. co.....
" senegæ.....aa fl. ℥ 1½.
Mix. Dose: a teaspoonful, for children.

Dr. Ackermann.

Letzerich's Treatment of Diphtheria by Benzoate of Soda.

℞ Sodæ benzoat, pur..... ʒ j.
Aquæ destillat.....
Aquæ menth. pip.....aa ℥ j.
Syrup cort. aurantii..... ʒ ij.
M. For an infant under one year old, one-half tablespoonful every hour.

For Hemorrhages.

℞ Acidi gallici..... ʒ j.
Glycerini..... ℥ ss.
Aquæ destillat..... ℥ vj.
M. Sig.—Two tablespoonfuls three times a day.

Farquharson.

In Mitral Regurgitation and in Dilatation of the Cavities of the Heart.

℞ Ferri redacti.....
Quiniæ sulphat.....
Pulv. digitalis.....aa ℈ j.
Pulv. scillæ.....gr. x.
M. Ft. pil. No. xx.
Sig.—One pill three or four times a day. *Bartholow.*

This prescription is especially serviceable in those cases of dilatation accompanied by cough, difficult breathing and general dropsy, and is of more value in such cases than so called cardiac sedatives and diuretics.

In Chronic Heart Disease.

℞ Pulv. digital.....gr. xxx.
Fer. sulph. exsiccat.....gr. xv.
Pulv. capsici.....gr. xl.
Pil. al. et myrrh..... ℥ ij.
M. Ft. pil. No. lx.
Sig.—One twice a day. *Fothergill.*

This is indicated in the following condition so frequent in heart disease, viz.: Where gastric catarrh, copious eructations of wind and inactivity of the bowels are associated with the cardiac debility.

Mixture of Digitalis and Iron for Cardiac Weakness with Dilatation of the Ventricles.

℞ Tr. ferri perchlorid..... ʒ ij.
Syr. zingiberis..... ʒ vj.
Inf. digitalis..... ℥ v.
M. Tablespoonful three times daily.

Bell's Gargle.

℞ Sodii biboratis.. ..... ʒ 2.
Fermenti.....
Mellis.....aa ℥ ½.
Aquæ q. s. ad.....fl. ℥ 8.
Mix.

In Cardiac Debility with Scanty Flow of Urine.

℞ Tinct. digitalis.....♏x.
Sp. æther nit..... ʒ ss.
Inf. buchu..... ℥ j.
M. Sig.—Three times a day. *Fothergill.*

This is a pleasant combination and may have citrate of potash added to it with advantage, or if there is much debility with the heart disease, or it is associated with atonic gout, the following will prove very efficacious:

℞ Pot. bicarb.....gr. x.
Fer. am. cit..... ..gr. v.
Tinct. digitalis.....♏x.
Inf. buchu ..... ℥ j.
Sig.—Three times a day, followed by draught of water.

Influenza.

℞ Ac. phos. dil.....M. xv.
Spt. chloroformi.....M. xxv.
Syr. scillæ..... ʒ j.
Aquæ.....ad. ℥ j.
M. Sig.—Four or six times every twenty-four hours.
Fothergill.

This will form an efficient expectorant, free from depressant effects in epidemic catarrh. In the convalescent stage where a tonic is also desired the following is very acceptable:

℞ Acid nit. dil.....M. x.
Sp. chloroformi..... ʒ ss.
Inf. cinchonæ..... ℥ j.

Sig.—Three or four times a day. *Fothergill.*

In Ulcerated Mouth or Gums or Mercurial Salivation.

℞ Potass. chloratis..... ʒ j.
Aquæ..... ℥ vi.

Two tablespoonfuls three times a day as a gargle.
Farquharson.

Acute Pleuritis.

℞ Tinct. aconit. rad..... ʒ ij.
Tinct. opii deodorat..... ʒ vj.

M. Sig.—Eight drops in water every hour or two.
Bartholow.

In acute pleuritis, before the stage of effusion has set in, no more effective combination will be found than the above.

Enlarged Heart Without Valvular Lesion.

℞ Ext. ergotæ fl..... ℥ iijss.
Tr. digitalis..... ℥ ss.

M. Sig.—A teaspoonful three times a day.
Bartholow

A Stimulating Expectorant.

℞ Am. carbonat.....gr. v.
Tinct. nuc. vom.....m. x.
Tinct scillæ..... ʒ ss.
Inf. serpentar..... ℥ j.
M. Sig.—Three times a day. *Fothergill.*

In those cases in which chronic bronchitis is associated with emphysema, or in the second stage of acute bronchitis, where the heart is severely taxed, this combination of remedies will strengthen the over-taxed heart and clear out the air passages.

Acute Catarrh.

℞ Antim. et potass. tart.....gr. ss.
Morphiæ acetat.....gr. ss.
Aquæ..... ℥ ij.
M. Sig.—A tablespoonful every hour or so.
Dr. Barthelow.

This is advised in the first stage of catarrh when the mucous membrane is dry and swollen.

Hæmoptysis.

℞ Plumbi acetat.....℈ ij.
Pulv. digital .....℈ j.
Pulv. opii.....gr. x.
M. Ft. pil. No. xx.
Sig.—One every four hours.
Dr. Roberts Bartholow.

Caseous Pneumonia.

℞ Inf. digital..... ℥ iv.
Plumbi acetat.... ..... ℈ j.
Tinct. opii..... ʒ j.
M. Sig.—A teaspoonful twice a day.

Dr. Oppolzer.

Coryza.

℞ Pulveris aluminis.....gr. v-xxx.
Aquæ..... ℥ j.
M. Sig.—To be drawn into the nostrils three or four times a day in the form of spray.

J. Solis Cohen, M.D.

This spray will diminish the profuse secretion and destroy the fœtor incident to coryza.

Expectorant Mixture for Children.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Syr. senegæ.....
" pruni virg.....
" acaciæ.....aa fl. ℥ i.
M. Dose: a teaspoonful. *Dr. Holgate.*

Expectorant Mixture.

Insane Asylum, Blackwell's Island, N. Y.

℞ Spts. etheris co..... } Equal parts of each.
Syr. ipecac..... }
Tinct. opii camph..... }
Aquæ..... }
M. Dose: a teaspoonful.

Cough Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Syr. tolut.....
Syr. pruni. virg.....
Tr. hyoscyami..... } aa fl. ℥ j.
Spts. etheris com.....
Aquæ.....

M. Sig.—Dose: a teaspoonful.

Prof. E. G. Janeway.

℞ Acid hydrocyan dil..... ʒ i.
Tinct. sanguinariæ..... ʒ iv.
Syr. senegæ..... ℥ ss.
Syr. tolutan..... ℥ ij.
Aquæ lauro-cerasi..... ʒ vij.

M. Sig.—One or two teaspoonfuls, according to age, every three or four hours. *Dr. Bartholow.*

This formula is most successful in treating the cases of cough by habit after the cessation of the whooping cough proper. It is also very useful in allaying the nervous cough of mothers which exists during the presence of cough in the household.

Expectorant Mixture.

Charity Hospital, Blackwell's Island, N. Y.

℞ Tinct. sanguinar..... fl. ʒ i
Tinct. opii camph.....
Syr. scillæ.....
Syr. tolut..... aa fl. ʒ ss
Aquæ q. s. ad..... fl. ℥ ss

Mix. Dose: a teaspoonful.

Asthmatic Mixture.

Bellevue Hospital, N. Y.
and
Charity Hospital, Blackwell's Island, N. Y.

℞ Spiritus æther co.....
Liquor morph. sulph (U. S.).....aa fl. ℥ i

Mix. Dose: from one teaspoonful to a tablespoonful.

Mistura "Bronchi."

Outdoor Department of Bellevue Hospital, N. Y.

℞ Ammonii carbon.....grs. 10
Syr. ipecac.....fl. ʒ i ½
Tinct. opii camph.....fl. ʒ i
Syr. pruni virg.....
Aquæ q. s. ad.....fl. ℥ ii

Mix. Dose: a teaspoonful, for children.

Dr. Beverly Robinson.

Carbonate of Ammonia Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Ammonii carbonat..... ʒ ½
Syr. senegæ.....fl. ʒ iv
" ipecac.....fl. ʒ ii
" tolut.....fl. ʒ iv
Ext. glycyrrh..... ʒ ½
Aquæ cinnam. q. s. ad.....fl. ℥ iv

Mix. Dose: a teaspoonful, for children.

Dr. Geo. H. Bosley.

Chloride of Ammonium Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Ammonii chloridi..... ʒ ½
Potassii chlorat.....grs. 40
Syr. senegæ.....fl. ʒ 4
" ipecac.....fl. ʒ 3
" tolut.....fl. ʒ 5
Ext. glycyrrhizæ..... ʒ 1
Aqua cinnam. q. s. ad.....fl. ℥ 4
Mix. Dose: a teaspoonful, for children.

Dr. Geo. H. Bosley.

Compound Licorice Mixture.

(*Brown Mixture.*)

℞ Pulv. ext. glycyrrh.....
" sacchari.....
" acaciæ.....aa ʒ 1
Tinct. opii camph.....fl. ℥ ½
Vini antimonii.....fl. ʒ 2
Spts. etheris nit.....fl. ʒ 1
Aquæ q. s. ad.....fl. ℥ 4
Mix. Dose: a teaspoonful. Shake before using.

Stokes' Expectorant Mixture.

℞ Ammonii carbon.....grs. 32
Ext. senegæ fl.....
Ext. scillæ fl.....aa fl. ʒ 1
Tinct. opii camph.....fl. ʒ 6
Aquæ.....fl. ℥ ½
Syr. tolut. q. s. ad.....fl. ℥ 4
Mix. Dose: a teaspoonful.

Expectorant Emulsion.

Hart's Island Hospital, New York.

℞ Morph. sulph.....gr. ij.
Syr. scillæ.....
" ipecac.....aa fl. ℥ ij.
" tolut.....
" pruni virg.....aa fl. ℥ jss.
Tr. benz. comp.....
Tr. sanguinariæ.....aa fl. ℥ ss.
Aquæ.....fl. ℥ ij.

Mix. Dose: a teaspoonful.

Expectorant Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Syr. scillæ co.....
" ipecac.....aa fl. ℥ i
Syrupi.....fl. ℥ i

Mix. Dose: a teaspoonful, for children.

Dr. Swezey.

Sedative Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Acid. hydrocyan. dilut.....
Chloroformi purif.....aa fl. ʒ i
Tinct. hyoscyami.....
Syr. tolutani.....
Aquæ camphoræ.....
Mucil. acaciæ.....aa fl. ℥ i

Mix. Dose: a teaspoonful.

Dr. Katzenbach.

Licorice Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Ammonii chloridi.....
Ext. glycyrrhizæ.....aa ʒ 2
Tinct. opii camph.....fl. ʒ 2
Aquæ q. s. ad.....fl. ℥ 4

Mix. Dose: a teaspoonful. *Dr. Brekes.*

Hoffmann's Anodyne and Iodide of Potash.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Potass. iodidi.....ʒ 3
Tinct. tolut.....fl. ʒ 1
Ext. pruni virg. fl.....fl. ʒ 1
Syrupi.....fl. ℥ 1
Spts. ætheris co.....fl. ʒ 2
Aquæ.....fl. ℥ 1

Mix. Dose: a teaspoonful.

Prof. E. G. Janeway.

Nitrate of Potash Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Potass. nitrat.....gr. 1
Spts. æther. nit.....
Syr. ipecac.....aa fl. ʒ ½
Syr. pruni virg.....fl. ʒ 2
Aquæ q. s. ad.....fl. ℥ 1

Mix. Dose: a teaspoonful, for children.

Dr. Robinson.

Hydrocyanic Mixture.

Bellevue Hospital, New York.

℞ Potass. cyanidi.....
Morphiæ sulphat.....aa grs. iv.
Syr. tolut.....fl. ℥ iv.

Mix. Each ʒ contains ⅛ gr. each of potassium cyanide and morphia sulphate. Dose: a teaspoonful.

Hydrocyanic Mixture.

Charity Hospital, Blackwell's Island, N. Y.

℞ Potass. cyanidi.....grs. 2
Vini antimonii.....fl. ʒ 2
Syr. tolut.....
Mucil. acaciæ.....aa fl. ℥ ½
Aquæ q. s. ad.....fl. ℥ i

Mix. Dose: a teaspoonful.

Chlorate of Potash Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Potass. chlorat..... ʒ i
Ext. glycyrrhizæ..... ʒ ½
Ammonii chloridi..... ʒ i
Aquæ.....fl. ℥ 4

Mix. Dose: a teaspoonful. *Dr. Geo. G. Wheelock.*

Stimulating Expectorant.

℞ Ammon. carbonat.....gr. xl.
Tinct. scillæ..... ʒ ij.
Syr. tolutani..... ʒ iij.
Infus. senegæ..... ℥ vij.

Mis. Ft. mistura et sig. two tablespoonfuls every four hours. *Dr. Farquharson.*

Cyanide and Bromide of Potash Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Potass. bromidi..... ʒ 4
Potass. cyanidi.....grs. 4
Syr. pruni virgin.....fl. ℥ 4
Mix. Dose: a teaspoonful.

Pneumonia.

℞ Ammon. carb..... .....grs. lxxx.
Mucilag.....
Aquæ .....āā fl. ℥ i.
Spts. vini gallici.....fl. ℥ ii.
M. Sig.—A tablespoonful three times daily.

When indicated, this is alternated with the following:

℞ Syrup scillæ.co..... ʒ iii.
Vini ipecac..... ʒ iss.
Morph. sulph.....gr. ss.
Aquæ.. .... ℥ iss.
Syrupi q. s. ad..... ℥ iv.
M. Sig.—A tablespoonful three times daily.

Where the dyspnœa becomes severe, with signs of increasing exhaustion, feeble heart action, etc., the amount of stimulants is increased to ℥ xii daily, and in addition:

℞ Infus. digitalis..... ʒ ii.
Sig.—Dose, every three hours.

Where dulness is present over more or less of the lung, ammonium muriate, grs. xx. three times daily is administered. *Dr. W. H. Thomson.*

Hoffmann's Anodyne and Iodide of Potash.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Ammonii carbon.....grs. 50
Potass. iodidi..... ʒ ʒ
Syr. pruni virg.....
Spts. æther. co.....aa fl. ℥ 1½
Mix. Dose: a teaspoonful.

Dr. Katzenbach.

Hydrothorax.

℞ Potass. carbon.....grs. xx.
Aquæ..... ℥ ss.
M. et add
Succi limonis..... ..... ℥ ss.
Sig.—Dose, every two hours.

With this also give:
Infus. digitalis..... ʒ j-ij.
three times daily. *A. Clark.*

Counter-Irritant in Pleurisy.

℞ Iodinii..... ʒ j.
Potass. iodidi..... ʒ ij.
Aq. destil.....
Alcohol .....āā ℥ ij.
M. *W. H. Draper.*

℞ Iodinii..... ℥ ijss.
Potass. iodidi..... ℥ ss.
Alcoholis..... ℥ xij.
Aquæ..... ℥ iv.
M. *F. Delafield.*

Dyspnœa of Valvular Disease.

℞ Liq. Morph. U. S. P. . . . . .
Spts. ether. co. . . . . āā ℥ i.
M. Dose, a teaspoonful. *A. Flint.*

Diuretic in Pleurisy.

℞ Potas. acetat. . . . . ʒ ss
Infus. digitalis. . . . . ℥ ss.
M. Sig.—Dose, every four hours. *A. Flint.*

Diuretic Pill in Pleurisy.

℞ Pulv. scillæ. . . . . .
Pulv. digitalis . . . . . .
Massa hydrarg. . . . . āā gr. j.
M. Ft. pil. *W. H. Draper.*

Acute Catarrhal Bronchitis.

℞ Ammon. chlorid. . . . . ʒ ij.
Potass. chlorat. . . . . ʒ j.
Aq. Cinnam. . . . . ℥ iij.
Syr. senegæ. . . . . ℥ ss.
Spts. ether nit. . . . . ℥ ss.
Ext. glycyrrh. . . . . ʒ iss.
M. Sig.—Tablespoonful every two hours.
J. R. Leaming.

Emphysema.

℞ Potass. iodidi. . . . . ʒ iij.
Ammon. carbon. . . . . grs. 50
Syr. pruni virg. . . . . .
Spts. ether. co. . . . . āā ℥ iss.
M. Dose, a teaspoonful. *A. Flint.*

Acute Pleurisy.

At the onset of the attack:—

℞ Tr. aconit. rad.....gtt. xx.
Tr. opii camph.....
Spts. etheris nit.....āā ℥ ss.
Liq. ammon. acet..... ℥ v.

M. Sig.—A tablespoonful every three hours.

J. H. Ripley.

Bronchitis.

℞ Potass. iodidi..... ʒ iij.
Tr. tolutan..... ʒ j.
Ext. pruni virg. fl..... ʒ j.
Syrupi..... ℥ j.
Spts. ether. co..... ℥ ij.
Aquæ..... ℥ j.

M. Dose, a teaspoonful. *E. G. Janeway.*

In Ulceration of the Larynx.

℞ Glycerini acidi carbolici.....
Succi conii .....āā ℥ j.

Misce. A teaspoonful to be put into a pint of water at the temperature of 170°, and the steam inhaled for fifteen minutes twice or thrice a day.

Dr. Dobell.

In Hoarseness from Weakness of Larynx.

℞ Acidi benzoici.....gr. ss.
Pastæ ribis rubri (red currant paste)...gr. x.

M. fiat trochiscus. One every hour or two.

Dr. M. Mackenzie.

In Cardiac Debility.

℞ Liq. strychniæ..... ♏ v.
Tinct. digitalis..... ♏ x.
Tinct. ferri perchlor..... ♏ x.
Aquæ vel inf. quassiæ..... ℥ j.

Misce. To be taken three times a day.

Dr. J. M. Fothergill.

Anti-Rheumatic in Pericarditis.

℞ Potass. iodidi..... ʒ v.
Vin. colchici sem..... ℥ i.
Tinct. cimicifug. rac..... ℥ ii.
Tinct. stramon..... ℥ ss.
Tinct. opii camph..... ℥ iss.

M. Dose: a teaspoonful three times daily.

Dr. W. H. Draper.

In Angina Pectoris.

℞ Liq. arsenicalis..... ♏ v.
Aquæ..... ℥ j.

Misce. To be taken three times a day.

Dr. F. E. Anstie.

In Hæmoptysis.

℞ Acidi tannici..... ʒ ss.
Conf. rosæ, q. s.....

Fiant pilulæ, xviij. Take one every four hours.

Cottereau.

In Ulcerated Throat, Cancrum Oris, etc.

℞ Cupri sulphatis.....gr. v.
Oxymellis..... ℥ ss.
Misce. Apply it with a camel-hair pencil.'

Mr. Brande.

In Consumptive Cough.

℞ Mucil. acaciæ.....f ℥ j.
Aquæ destil.....f ℥ viss.
Syr. tolutani.....f ℥ ss.
Acidi hydrocy. dil.....♏ xij.
Misce. A tablespoonful every three hours.

Mr. S. G. Morton.

In Cardiac Debility.

℞ Tinct. belladonnæ.....♏ xx.
Tinct. nucis vom.....♏ x.
Aquæ camph..... ℥ j.
Misce. To be taken three times a day.

Dr. J. M. Fothergill.

To Quiet Common or Spasmodic Cough.

℞ Ext. conii.....gr. v.
Mist. amygdalæ.....f ʒ xiv.
Syr. papaveris.....f ʒ ij.
Misce. Sumat cochleare unum medium bis vel ter die.

Mr. Brande.

In Advanced Heart Disease.

℞ Gambogiæ.....gr. j.
Pulv. jalapæ co..... ʒ ij.
Fiat pulvis.

Dr. J. M. Fothergill.

In Cardiac Debility.

℞ Tinct. colchici sem.....♏ x.
Tinct. digitalis.....♏ x.
Sp. etheris nit.....♏ xxv.
Inf. buchu.....℥ j.
Misce. To be taken three times a day.

Dr. J. M. Fothergill.

In Acute Congestion of the Pharynx and Larynx.

℞ Tinct. benzoini co.....ʒ j ad ʒ ij.

To be added to a pint of water at 150° F., and inhaled for five or eight minutes every four or six hours from a suitable inhaler. *Dr. Morell Mackenzie.*

In Chronic Heart Disease.

℞ Pulv. digitalis.....gr. ½—j.
Ferri sulph.....gr. j.
Pulv. capsici.....gr. ¼.
Ext. gentianæ.....gr. ij.
M. f. pilula. Take three a day. *Dr. Fothergill.*

In Simple Cardiac Debility.

℞ Tinct. digitalis.....♏ x.
Sp. chloroform.....♏ xxv.
Inf. buchu.....℥ j.
Misce. To be taken three times a day.

Dr. J. M. Fothergill.

In Chronic Pleuritis.

℞ Pulv. digitalis.....
Quin. sulph..... .....aa gr. j.
Confec. rosæ, q. s.....
Fiat pilula. To be taken three times a day.

Niemeyer.

In Cardiac Debility.

℞ Ferri tartarat.....gr. v.
Potass. bicarb.....gr. v.
Tinct. digitalis.....♏ x.
Inf. calumbæ..... ℥ j.
Misce. To be taken three times a day.

Dr. J. M. Fothergill.

In Tonsillitis.

℞ Resinæ guaiaci.....gr. iij.
Pastæ ribis rubri (red currant paste)..gr. vij.
M. f. trochiscus. [One to be sucked every hour or two.] *Dr. M. Mackenzie.*

In Ozæna.

℞ Hydrarg. iodidi rubri.....
Hydrarg. subchlor.....aa gr. xij.
Sacch. alb..... ℥ ss.
Misce. To be used as an errhine. *Niemeyer.*

In Acute Bronchial Catarrh.

℞ Ext. glycyrrhizæ..... ℥ ij.
Liq. ammon. anisati..... ℥ ij.
Aquæ fœniculi..... ℥ vj.
Misce. To be taken in teaspoonfuls. *Niemeyer.*

In Bronchocele.

℞ Tinct. iodi..... ℥ j.
Inject ♏xxx into the substance of the gland once a week for the first two or three weeks, and after, once a fortnight as long as necessary. Give iodide of potassium internally. *Dr. Morell Mackenzie.*

In Pleuritis.

℞ Iodi ..... ʒ ss.
Potass. iodidi ..... ʒ ss.
Aquæ destil ..... ℥ ij.
Fiat lotio. To be applied to affected side.

Niemeyer.

In Ozæna.

℞ Iodini ..... gr. ij—iv.
Potass. iodidi ..... gr. iv—viij.
Aquæ ..... ℥ vj.
Fiat injectio. *Niemeyer.*

An Emetic in Nocturnal Dyspnœa of Laryngitis.

℞ Pulv. ipecac ..... ☉ j.
Antimon. tartar ..... gr. j.
Oxymel. scillæ ..... ʒ iij.
Aquæ ..... ℥ iss.
Misce. A teaspoonful every ten minutes.

Niemeyer.

In Putrid Sorethroat, Scarlatina, etc.

℞ Tinct. myrrhæ .....
Aceti ..... ana ℥ ij.
Mellis ..... ℥ j.
Inf. serpentariæ ..... Oiiss
Misce. Fiat gargarisma. *Dr. Fothergill.*

In Relaxation of Pharynx.

℞ Ex. krameriæ ..... gr. iij.
Pastæ ribis rubri (red currant paste) .. gr. vij.
M. Fiat trochiscus. One every two or three hours.

Dr. M. Mackenzie.

A Stimulant in Congestion of the Larynx.

℞ Ol. pini sylvest..... ʒ ij— ʒ iij.
Magnes. carb. levis.....gr. lx.
Aquæ, ad..... fl. ℥ iij.

M. A teaspoonful in a pint of water, at 150° F., to be inhaled night and morning for ten minutes. (An increased stimulant character may be given to this inhalation by the addition of twenty grains of camphor to the mixture.) *Dr. Morell Mackenzie.*

In Acute Bronchial Catarrh.

℞ Liq. ammon. anisat..... ʒ ss.
Inf. senegæ..... ʒ ij-iv.
Aquæ..... ℥ vj.

Misce. Two tablespoonfuls three times a day.
Niemeyer.

In Aphthæ of the Mouth.

℞ Sodæ sulphitis..... ʒ j.
Aquæ..... ℥ j.

Misce. Ut fiat lotio. *Sir W. Jenner.*

In Dropsy of Emphysema.

℞ Aceti scillæ..... ℥ j.
Potass. carb. ad saturationem.....
Aquæ destil..... ℥ vj.

Misce. A tablespoonful every two hours.
Niemeyer.

Heart Stimulant.

℞ Nitroglycerini.....gr. $\frac{1}{100}$
Tr. digitalis.....♏ij.
Tr. strophanthi.....♏ij.
Tr. belladonnae.....♏$\frac{1}{4}$

M. Make into tablet or pill.

Sig.—One every three to four hours for *angina pectoris*, *palpitation*, *tobacco heart*, or *weak heart*.

Dr. DaCosta.

For Dilated Heart.

℞ Potass. iodid.....gr. v.
Ex. digitalis fl.....♏ij.
Ex. convallariæ majalis fl.....♏xx.

M. Sig.—For a dose repeated after each meal.

Delafield.

For Acute Cold.

℞ Quininæ sulphatis.....gr. ij.
Pulv. ipecac. et opii.....gr. ij.
Oleores. capsici.....gr. $\frac{1}{60}$
Tr. aconiti.....♏ss.

M. Ft. tablet.

Sig.—One every three hours for acute cold.

For Bronchitis.

℞ Terebene.....ʒj.

Sig.—Ten to fifteen drops on sugar every four hours in acute or chronic bronchial catarrh, especially with profuse muco-purulent expectoration.

The terebene may also be administered according to the following formula:

℞ Terebene..... ..... ʒ ijss.
Mucl. acacia..... .....
Aquæ..... aa ℥ ss.
Syr. zingiberi..... .. ..q. s. ad ℥ ij.
M. Sig.—Teaspoonful three times a day. *Nicholls.*

For Cough.

℞ Terpin hydratis..... ...gr. v.
Codeinæ.....gr. ¼
M. Ft. tablet.

Sig.—One every two to four hours for cough due to *laryngitis* or *bronchitis.* A very effective tablet for harrassing *night cough.*

For Cough.

℞ Heroin.....gr. $\frac{1}{24}$
Ft. tablet.

Sig —One tablet every three to four hours for cough or pains in the chest due to any cause, but especially *bronchitis* and *phthisis.* The dose of heroin ranges from one-twelfth to one-fiftieth of a grain. The former dose sometimes causes nausea; the latter is frequently sufficient to control cough, but is insufficient in many cases. Heroin may be satisfactorily combined with terpine as in the following prescription:

℞ Heroin.....gr. $\frac{1}{24}$.
Terpin. hydrat.....gr. iii.-v.
M. Ft. tablet.

Sig.—One every three to four hours. This is of especial value in the cough of elderly people.

For Epistaxis.

℞ Antipyrin.....gr. l.
Acid. tannic .....gr. j.
Pulv. sacchari.....gr. x.
M. Sig.—Apply topically. *Rendu.*

For Quinsy.

℞ Sodii salicylat ..... ʒ iij.
Syr. acaciæ..... ℥ ss.
Aq. cinnam.....ad ℥ iij.
M. Sig.—Dessertspoonful every three hours.
Easby.

For Dry Pharyngitis.

℞ Pilocarpinæ muriat.....gr. ij.
Aquæ.....
Glycerinæ.....aa ℥ j.
M. Sig.—Teaspoonful three times a day. *Sajous.*

For Sore Throat.

℞ Menthol.....gr. $\frac{1}{35}$
Ac. benzoici.....gr. $\frac{1}{12}$
Eucalyptol.....gr. $\frac{1}{16}$
Ol. Gaultheriæ.....gr. $\frac{1}{16}$
Cocainæ hydrochlor.....gr. $\frac{1}{280}$
Sacchari.....gr. v.
M. Sig.—Dissolve one in the mouth every hour or less.

This is an excellent stimulant and antiseptic tablet which does not *dry* the throat nor disturb the appetite. It is indicated in all forms of sore throat having a red and inflamed mucous membrane.

For Phthisis.

℞ Terpinol.....
Sodii benzoatis.....aa gr. xv.
M. Div. in capsulas No. x.
Sig.—A capsule every hour or two. To diminish the expectoration and remove its odor. *Rabow.*

Coryza—Rhinitis.

℞ Camphoræ.....gr. ss.
Ext. belladonnæ fl.....♏¼
Quininæ sulphatis.....gr. ss.
M. Ft. tablet.
Sig.—One tablet every hour for four doses, then every two to four hours. *Dr. Lincoln.*

This is a very popular tablet and is extensively used. The half strength tablets frequently employed must be administered very frequently to be effective. The following prescriptions are especially effective in the coryza accompanying *grip*, either with or without sore throat and cough:

℞ Quininæ sulphatis.....
Camphoræ.....
Ammonii muriatis.....aa gr. ss.
Pulv. opii.....
Ext. belladonnæ.....
Ext. aconiti.....aa gr. $\frac{1}{10}$
M. Ft. tablet.
Sig.—One every hour till the throat becomes dry, then every two to three hours. *Dr. Richards.*

℞ Quininæ sulphatis.....gr. $\frac{1}{8}$
Camphoræ.....gr. $\frac{1}{4}$
Morphinæ.....gr. $\frac{1}{80}$
Atropiæ sulphatis.....gr. $\frac{1}{600}$
Strychniæ sulphatis.....gr. $\frac{1}{240}$

M. Ft. tablet.
Sig —Administer like the preceding tablet.

For Otitis and Otorrhœa.

℞ Liq. hydrogenii peroxidi (10 vol.)..... ℥ iv.

Sig.—Syringe the ear carefully with one part solution to two of water, and when cleansed drop in a few drops of the above solution. *C. H. Burnett*

For Furunculosis of the Ear.

℞ Acid. carbol..... ..... ʒ ss.
Glycerinæ..... ..... ʒ ix.

M. Sig.—Drop a few drops into the ear two or three times daily, after cleansing. *Hartmann.*

For Earache.

℞ Chloral camphorat.....gr. v.
Glycerinæ.....gr. xxx.
Ol. amygdal. dulc..... .....gr. x.

M. Sig.—Apply a little on absorbent cotton and place in ear. *Journ. de Médecine.*

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

II.

DISEASES OF THE ABDOMINAL VISCERA.

In Lead Colic.

℞ Aluminis..... ℥ ij.
Acid sulph. dil..... ʒ j.
Syr. limonis..... ℥ j.
Aquæ..... ℥ iij.

M. Sig.—A tablespoonful every hour or two.

Bartholow.

Will relieve the violent pains of painter's colic.

Rum Stomach.

℞ Tinct. nucis vomicæ.....grs. v-xv.
Tinct. gentianæ co.....
Tinct. columbæ co.....aa ʒ j.

M. Sig.—Take before meals.

Dr. Alfred L. Loomis.

This is advised in those cases of rum stomach accompanied by gaseous distension of the bowels and stomach. We may also advantageously give a mercurial or aloetic purge.

Gastric Catarrh.

℞ Aluminis..... ..... ʒ ij.
Ext. gentian..... ʒ ss.

M. Ft. pil. No. xxx. Sig.—Two pills three times a day. *Dr. Bartholow.*

This formula is held to be most effective when there is vomiting of glairy mucus from the stomach.

West's Mixture.

℞ Ol. ricini.....fl. ʒ 2
Pulv. acaciæ.....
Pulv. sacchari.....aa ʒ 2
Tinct. opii.....m. 21
Aquæ cinnam. q. s. ad.....fl. ℥ 4

Mix. Dose: a teaspoonful, for children, in diarrhœa.

Opium, Rhubarb and Camphor.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Tinct. opii.....
Tinct. rhei arom.....
Spts. camphoræ.....aa fl. ʒ ½
Tinct. cardam co.....fl. ʒ 2
Aquæ anisi q. s. ad.....fl. ℥ 4

Mix. Dose: a tablespoonful, for children, in diarrhœa. *Dr. Swezey.*

Hot Drops.

Charity Hospital, Blackwell's Island, N. Y.

℞ Tinct. opii.....
Tinct. capsici.....
Spts. camphoræ.....
Spts. menthæ pip.....
} aa fl. ʒ 2
Aquæ.....fl. ℥ 1

Mix. Dose: a teaspoonful.

Cholera Morbus.

℞ Chloral hydratis..... ʒ iij.
Morphiæ sulphatis.....gr. iv.
Aquæ laurocerasi..... ℥ j.

M. Sig.—From fifteen to twenty minims injected hypodermically for cholera and cholera morbus.

Roberts Bartholow, M.D.

This injection may produce considerable burning pain and an indurated lump, but it is considered by Dr. Bartholow to be the most effective remedy known for relief of cholera, etc., the patient not infrequently being wrested from a condition of extreme danger.

Habitual Constipation.

℞ Ext. stillingiæ fl..... ʒ v.
Tinct. belladonnæ.....
Tinct. nucis vomicæ.....
Tinct. physostigmæ.....aa ʒ j.

M. Sig.—Twenty drops in water three times a day before meals. *Bartholow.*

Acute Dysentery.

℞ Cupri sulph.....gr. ss.
Magnes. sulph..... ℥ j.
Acid sulph. dil..... ʒ j.
Aquæ..... ℥ iv.

M. Sig.—A tablespoonful every four hours.

Dr. Roberts Bartholow.

Chronic Diarrhœa.

℞ Bismuthi subnitratis.....gr. v.
Morphiæ sulphat.....gr. $\frac{1}{12}$.

M. Sig.—This much two or three or more times daily. *Dr. Alonzo Clark.*

This formula will often afford relief in these cases, though the influence of proper diet and change of surroundings should also be considered in treatment.

Constipation.

℞ Tinct. nucis vomicæ.....
Tinct. belladonnæ.....
Tinct. physostigmæ.....aa ʒ ij.

M. Sig.—Thirty drops in water morning and evening. *Dr. Bartholow.*

Where constipation is due to torpor of the muscular layer of the intestine, combined with deficient secretion of the mucous membrane, this formula is often very serviceable, or the following may be taken:

℞ Ext. physostigmæ.....
Ext. belladonnæ.....
Ext. nucis vomicæ.....aa gr. v.

M. Ft. pil. No. 10. Sig.—One pill at bed hour. *Bartholow.*

Diarrhœa.

℞ Tinct. colombæ..... ʒ xv.
Tinct. opii deodor..... ʒ j.

M. Sig.—A teaspoonful in a wineglassful of water, before meals. *Bartholow.*

This formula is best adapted to a condition such as the following, viz.:

Soon after taking food, *pain* referable to small intestine, nausea, loose evacuations containing undigested aliments and followed by weakness and depression.

In First Stages of Cirrhosis.

℞ Ammon. iodid..... ʒ j.
Liq. potass. arsenit..... ʒ ss.
Tinct. colombæ..... ℥ ss.
Aquæ..... ℥ jss.

M. Sig.—A teaspoonful three times a day before meals. *Bartholow.*

Diarrhœa Mixture.

℞ Tinct. opii.....
Tinct. capsici.....
Tinct. rhei arom..... } aa p. e.
Spts. menthæ pip.....
Spts. camphoræ .....

Mix. Dose: 20–40 min. *Dr. Ruschenberger.*

Compound Aloes Mixture.

℞ Aloes socotr..... ℥ i
Sodii bicarb..... ℥ i½
Glycerine.....
Spts. lavand co .....aa fl. ℥ 2.
Ol. menthæ pip.....m. 25
Aquæ.....O i

Mix. Dose: one to two tablespoonfuls as a purgative; one to two teaspoonfuls as a stomachic and tonic.

Diarrhœa Mixture.

Hart's Island Hospital, N. Y.

℞ Tinct. capsici.....fl. ℥ i
Tinct. catechu.....
Tinct. kino.....
Tinct. krameriæ.....aa fl. ℥ 4
Tinct. opii.....fl. ℥ 3
Spts. menth. pip.....fl. ℥ 2
Spts. camphoræ.....
Aquæ.....aa fl. ℥ 4
Mix. Dose: 30–60 minims.

Diuretic Mixture.

℞ Spiritus juniperi.....℥ ss.
Potass. acetat.....ʒ jss.
Spiritus ætheris nit.....℥ j.
Decoc. scoparii (br).....℥ viij.
Dose : two tablespoonfuls three times a day.

Hyperæmia of the Kidney.

In addition to general hygienic measures, Prof. Montrose Pallen, M.D., advises the use of vaseline inunctions and sponging the surface with hot water and alcohol. The use of the hot air or Turkish bath, and the most easily digested and highly nutritious food, such as milk and kumyss. An aperient is to be given and a uniform temperature maintained by keeping the patient in bed. Massage is advantageous, stimulating the circulation. Should the symptoms of eclampsia come on, we must have recourse to chloroform and bleeding.

Gastralgia.

℞ Argenti oxidi.....grs. v.
Ext. hyoscyami.....grs. v.

M. Ft. pil. No. x. Sig.—One pill three times a day before meals. *Bartholow.*

This formula is indicated in cases where there is pain after taking food, lasting for an hour or more, burning pain with pyrosis coming on after the completion of the stage of stomach digestion, eructations of food, with sour and acrid matters.

Anti-Emetic Mixture.

Charity Hospital, Blackwell's Island, N. Y.

℞ Creasoti.....m. 12
Acid hydrocyan. dil.....m. 30
Pulv. acaciæ.....
" sacchari.....aa ʒ 6
Aquæ q. s. ad.....fl. ℥ 2

Mix. Dose: a teaspoonful.

Castor Oil Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Olei ricini.....fl. ʒ 4
Mucil. acaciæ.....ʒ 4
Tinct. opii.....fl. ʒ 2
Tinct. rhei. arom.....fl. ʒ 4
Aquæ menthæ pip. q. s. ad.....fl. ℥ 4

Mix. Dose: a teaspoonful, for children, in diarrhœa. *Dr. Bosley.*

Hope's Mixture.

℞ Acid nitrici.....m. 8
Tinct. opii.....m. 40
Aquæ camphoræ.....fl. ℥ 8
Mix. Dose: a tablespoonful. In dysentery.

Rhubarb and Soda.

Outdoor Department of Bellevue Hosptal, N. Y.

℞ Sodii bicarb ..... ʒ 1
Extr. rhei. fl.....
Spts. menthæ pip.....aa fl. ʒ 1
Aquæ q. s. ad..... ....fl. ℥ 4
Mix. Dose: a tablespoonful.

Compound Rhubarb Mixture.

1 fl. ʒ contains.

℞ Ext. rhei. fl.....m. 256.....m. 1
Ext. ipecac fl.....m. 51.....m. ⅕
Sodii bicarb.....grs. 512.... .grs. 2
Glycerinæ.....fl. ℥ 12.....fl. ʒ ⅖
Aq. menthæ pip.....O 2.....fl. ʒ ⅗
Mix. Dose: ½ to 1 teaspoonful, two or three times a day, for children. *Dr. Squibb.*

Bicarbonate of Soda.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Sodii bicarb..... ℥ 1
Tinct. zingiber.....fl. ʒ 2
Tinct. gent. co.....fl. ℥ ʒ 1
Aquæ.....fl. ℥ ʒ 5
Mix. Dose: two teaspoonfuls.

Rhubarb and Soda.

Bellevue Hospital, New York.

℞ Sodii bicarbon..... ʒ i
Pulv. rhei..... ℥ ½
Spts. menthæ pip.....fl. ʒ ii
Aquæ q. s. ad.....fl. ℥ iv
Mix. Dose: a tablespoonful.

Rhubarb and Lime.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Tinct. opii camph .....
Syr. rhei. arom.....aa fl. ℥ ½
Aquæ calcis.....fl. ℥ ii
Mix. Dose: a teaspoonful, for children, in diarrhœa. *Dr. Ackermann.*

In Gastralgia and Irritative Dyspepsia.

℞ Bismuthi subcarb..... ʒ iij.
Morph. sulph.....gr. j.
Pulv. aromat..... ʒ j.
Ft. pulv. No. xii. Sig.—A powder in milk before each meal. *Dr. Bartholow.*

The above will quiet the existing irritability and relieve the pain in these cases.

In Acute and Chronic Dysentery.

℞ Plumb. acetat.....grs. iv.
Morphiæ acetat.....gr. ss.
Aq. fervid..... ℥ j.
M. Sig.—Use as an enema.

Dr. Roberts Bartholow.

Chronic Interstitial Nephritis.

℞ Tinct. ferri chloridi.....m. x.
Syrupi limonis.....m. l.
Aquæ.....℥ iij.

M. Sig.—To be taken thrice daily in a wineglassful of water. *Dr. Da Costa.*

Patients in addition to the above means of medication, should be treated by a nourishing diet, a little wine is allowable, and as a matter of course, hygiene should be strictly attended to.

Uræmia.

℞ Acid benzoici.....gr. xx.
Syrupi tolutani.....℥ j.

Sig.—To be given every third hour, largely diluted with water. *Dr. Da Costa.*

Benzoic acid has the effect of preventing the accumulation of the urinary salts in the blood, and thus exercising a favorable influence over the course of the disease. Dr. Da Costa advises also warm bathing (and if patient is very weak a vapor bath may be given him in bed by means of hot bricks wrapped in wet towels) and an efficient purge.

For Sluggish Liver.

℞ Acidi nitro-muriat. dil.....℥ ii.
Succi (extract) taraxaci.....℥ ss.
Spiritus chloroformi.....℥ jss.
Aquæ.....ad. ℥ viij.

Sig.— ℥ j. three times daily. *Dr. Farquharson.*

Diarrhœa of Phthisis.

℞ Resin terebinth.....gr. iii.
Argenti nitrat.....
Opii.....aa gr. ¼

M. Sig.—One pill when needed.

This formula is used in these cases at Roosevelt Hospital with uniform good results, the diet being milk boiled with mutton suet until it is as thick as cream.

Squibb's Cholera Mixture.

℞ Tinct. opii.....
" capsici.....
Spts. camphoræ.....aa fl. ℥ i
Chloroformi.....fl. ʒ ss
Alcoholis q. s. ad.....fl. ℥ v

Mix. Dose: 20–40 minims.

Pulv. Bismuthi Cretæ et Opii.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Bismuth. sub-nitr.....gr. 30
Cretæ præcipit.....gr. 30
Pulv. opii.....gr. i

Mix. Divide into ten powders. For children in diarrhœa. *Dr. J. Lewis Smith.*

Cole's Laxative Pills.

℞ Extr. colocynth. co.....gr. 30
Hydrarg. chlor. mit.....gr. 10
Resin. podophylli.....gr. i

Mix. Divide into ten pills.

Pillulæ Aperientes.

℞ Ext. aloes.....gr. 20
Pulv. rhei.....gr. 10
Ext. nucis vom.....gr. 5
Ext. taraxaci.....gr. 30

Mix. Divide into twenty pills. *Dr. Lusk.*

Pillulæ Aloes et Fellis.

99th St. Reception Hospital, N. Y.

℞ Extr. aloes .....gr. 30
Fellis bovis purif.....gr. 20
Resin. podophylli.....gr. 2½

Mix. Divide into ten pills. Dose: One pill at night in chronic constipation; one pill night and morning in acute constipation.

Laxative Pills.

99th St. Reception Hospital, N. Y.

℞ Pulv. rhei.....
Pulv. aloes.....aa gr. 15
Extr. bellad.....
Extr. nucis vom.....
Resin. podophylli.....aa gr. 3
Olei caryophylli.....gtt. 5

Mix. Divide into twelve pills. Dose: One pill mornings and evenings.

Squibb's Laxative Pills.

℞ Resinæ podophylli.....gr. 36
Extr. bellad. alcoh .....gr. 18
[*or*, Extr. hyoscyam. alc.....gr. 144]
Pulv. capsici .....gr. 144
Pulv. sacch. lactis.....gr. 144
Pulv. acaciæ .....gr. 36
Glycerinæ .....m. 40
Syrupi.....q. s.

Mix. Divide into 144 pills. Dry them by exposure to the ordinary temperature, until just hard enough to retain their form. Then put them into a well-stoppered bottle.

"Vegetable Cathartic Pills."

℞ Extr. colocynth. co .....gr. 36
Resin. podophylli .....gr. 9
Resin. leptandræ .....gr. 3
Pulv. jalapæ .....gr. 6
Pulv. aloes soc.....gr. 12
Extr. hyoscyam.....gr. 6
Olei menthæ pip.....gtt. 5

Mix. Divide into twenty-four pills

Liquid Pepsin.

℞ Pepsini. .....gr. 64
Aquæ.....fl. ℥ 2½
Acidi hydrochlorici.....fl. ½
Glycerinæ.....fl. ℥ 1½

Mix and filter.

Pulv. Bismuthi et Doveri.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Bismuthi sub-nitr.....gr. 4
Pulv. ipecac co.....gr. 1

Mix. One dose for children, in diarrhœa.

Pulvis Bismuthi Co.

Outdoor Department of Bellevue Hospital.

℞ Bismuthi sub-nitr.....
Sodii bicarb.....
Pulv. sacchari..... } aa p. e.
Pulv. acaciæ.....
Pulv. zingiber.....

Mix. Dose: a tablespoonful, for adults, in dyspepsia. *Dr. Wheelock.*

Pulv. Bismuthi et Pepsini.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Bismuthi sub-nitr.....
Pepsini .....aa gr. 3

Mix. One dose, for children. *Dr. Swezey.*

Pulvis Glycerrhizæ Co.

℞ Pulv. sennæ.....part 2
Pulv. glycyrrh. rad..... " 2
Pulv. foeniculi..... " 1
Sulphuris loti..... " 1
Pulv. sacchari..... " 6

Mix. Dose: 30 to 60 grains; an excellent and pleasant laxative.

Pulv. Bismuthi et Pepsini.

Infants' Hospital, Randall's Island, N. Y.

℞ Bismuthi sub-carb.....
Pepsini.....aa gr. 2
Mix. One dose for children.

Diarrhœa Powder.

Infants' Hospital, Randall's Island, N. Y.

℞ Bismuthi sub-carb.....gr. 2
Acid tannici.....gr. 1
Pulv. ipecac co.....gr. ¼
Mix. One dose, for children, in diarrhœa.

The following formula I have sometimes used to fulfil the two-fold indication of relieving nausea and restraining the bowels in the acute stages of gastro-enteritis.

℞ Bismuth. ammon. cit..... ʒ ss.
Acid carbolic.....gtt. ij.
Liqr. atropiæ.....gtt. ij.
Tr. menthæ.....gtt. viij.
Tr. opii. deoderat.....gtt. x.
Syrup. acaciæ..... ℥ ij.
Mix. Sig.—A teaspoonful every two or three hours.

When the more violent symptoms have abated I frequently prescribe the following to advantage:

℞ Bismuth sub. nitrat..... ʒ i.
Lactopeptine..... ʒ ss.
Pulv. cretæ co. c. opii ..... ℈ ss.
M. Ft. chart No. x. Sig.—One three or four times daily.

When the evacuations are frequent, small, slimy or bloody, in other words, dysenteric, I find occasional use of a castor oil mixture to very great advantage, thus:

℞ Ol. ricini..... ʒ i.
Tr. opii. deoderat.....gtt. vi.
Syrup. acaciæ..... ℥ i.
Tr. menthæ.....gtt. ij.

Mix. Sig.—A teaspoonful every two hours till the stools assume a healthier character, after which the bismuth and lactopeptin powders may be resumed, with or without the comp. chalk and opium, as the condition of the bowels may indicate.

A. N. Talley, M.D., Columbia, S. C.

For Debility and Want of Appetite.

℞ Quiniæ sulph.....gr. viij.
Acid nit. dil..... ʒ ss.
Tinct. aurantii..... ℥ ss.
Syrupi aurantii..... ℥ j.
Aquæ.....ad. ℥ viij.

Dose: two tablespoonfuls three times a day.

Farquharson.

Diarrhœal Mixture.

℞ Tinct. catechu ..... ʒ vj.
Pulver. cret. aromat..... ʒ ij.
Tinct. opii..... ʒ jj.
Mucilaginis..... ℥ j.
Aquæ cinnamonis.....ad. ℥ vj.

Take a tablespoonful after each liquid dejection.

Farquharson.

Good Purgative Formula.

℞ Hydrarg. subchlor.....gr. xij.
Mannæ.....gr. vj.
Pulv. tragacanthæ comp.....gr. vj.
M. Divide into six pills and take two as required.
Farquharson.

Chronic Gastric Catarrh.

℞ Argenti nitrat.....gr. xv.
Aq. destillat q. s. ad. solv.....
Ext. belladon.....gr. x.
Ol. carophylli.....gtt. x.
Rad. gent. pulv.....
Ext. gentianæ aa q. s. ut pill No. 60.....
Sig.—One three times a day. *Bartholow.*

Chronic Diarrhœa and Dysentery.

℞ Cupri sulphat.....gr. j.
Morphiæ sulphat.....gr. j.
Quiniæ sulphat.....gr. xxiv.
M. ft. pil. No. xij.
Sig.—One pill three times a day. *Bartholow.*

Dr. Bartholow says of the above prescription: "Of all the metallic astringents employed for this purpose, I know of none better or more effective than sulphate of copper. It is specially indicated where there are colic-pains, tenesmus, and the stools contain mucus streaked with blood."

In Gastric Ulcer.

℞ Bismuthi subnitrat..... ʒ ij.
Morphiæ sulphat.....gr. j.
M. ft. pulv. No. vj.
: Sig.—One three times a day in milk.

Bartholow.

OR

℞ Bismuthi subnitrat..... ʒ ij.
Acid. hydrocyanici dil..... ʒ ij.
Mucilag. acaciæ.....
Aquæ menth. piperit.....aa ℥ ij.
M. Sig.—Tablespoonful three times a day.

Bartholow.

These formulæ will relieve the pain and vomiting of gastric ulcer, which is often so persistent.

Laxative.

℞ Magnes. sulph..... ℥ ij.
Ferri sulph.....gr. xxiv.
Acid. sulphur. dil..... ʒ ij.
Infus. quassiæ..... ℥ viij.
Ft. mistura. Take a tablespoonful when going to bed.

Farquharson.

A Good Dinner Pill.

℞ Ferri sulph.....
Ext. nucis vomicæ.....aa gr. ss.
Ext. aloes barb.....gr. iij.
Ft. pil. To be taken before dinner.

Farquharson.

In Gastric Irritability.

℞ Bismuthi trisnitrat.....gr. x.
Pot, bicarb.....gr. v.
Mist. acaciæ..... ʒ j.
Inf. columbæ..... ℥ j.
M. Sig.—Three times a day before food.

Fothergill.

In tuberculosis, in illy-nourished patients where gastric irritability interferes with assimilation and digestion, the above mixture will quiet irritability and thus promote assimilation and improve nutrition.

For Flatulent Dyspepsia.

℞ Sodii bicarb.....
Sacchari.....aa ʒ ij.
Spiritus ammon. aromat.....m. xl.
Aquæ menth. piperit.....ad. ℥ viij.

The above formula is known as soda mint, and is a very pleasant antacid combination.

Dose.—A tablespoonful after meals.

Farquharson.

It admits of the addition of tinct. nux vomica or syrup of rhubarb.

Flatulent Colic.

℞ Tinct. nucis vomicæ..... ʒ j.
Acidi nitro muriatici dil..... ʒ ij.
Spiritus chloroformi..... ʒ j.
Infus. gentianæ.....ad. ℥ vj.

Dose.—Tablespoonful three times daily after meals.

Farquharson.

In Atonic Dyspepsia.

℞ Inf. columbæ..... ℥ iv.
Liq. potassæ ..... ʒ ss.

M. Sig.—A dessert to a tablespoonful three times a day before meals. *Bartholow.*

Anthelmintic.

℞ Olei terebinthinæ..... ℥ ss.
Olei ricini..... ℥ j.

Take in one dose. *Farquharson.*

The above prescription is very effective against round worms and also tæniæ.

For Irritative Dyspepsia.

℞ Acidi hydrocyanici dil.....m. xxv.
Bismuth subnitratis..... ʒ ss.
Syr. aurantii..... ℥ j.
Gentian. infusi.....ad. ℥ viij.

Sig.—A teaspoonful three times a day before meals.
Dr. Farquharson.

In Flatulence.

℞ Acidi carbolici.....gr. j.
Pulv. rhei.....gr. iij.

Fiat pilula. (One every morning after breakfast.)
Dr. Habershon.

In Sarcina Ventriculi.

℞ Sodæ sulphitis.....gr. xxx—xl.
Inf. quassiæ..... ℥ iss.

Fiat haustus. To be taken three times a day.
Sir W. Jenner.

In Constipation.

℞ Sulphur. sublim..... ʒ ij.
Potass. acid tart..... ℥ ss.
Syr. limonis.....
Saccha. alb.....āā ℥ iij.

Fiat electuarium. A teaspoonful two or three times a day. *Niemeyer.*

In Nephritis (second stage).

℞ Tinct. ferri perchlor..... ʒ ij.
Sp. ætheris nit..... ʒ iv.
Inf. quass. ad..... ℥ vj.

Misce. A tablespoonful three times a day.
Dr. Grainger Stewart.

In Acute Nephritis.

℞ Inf. digitalis..... ℥ iss.
Sp. ether nit..... ʒ vj.
Syr. simplicis..... ℥ ss.
Aquæ, ad ..... ℥ vj.

Misce. A tablespoonful three times a day.
Dr. Grainger Stewart.

A Strong Purge.

℞ Ol. crotonis..... ♏ iij.
Pulv. acaciæ.....gr. iij.

Tere simul, et adde.
Syr. simp..... ℥ ss.
Inf. anthemidis..... ℥ iij.

M. In one or two doses. *Trousseau and Reveil.*

In Chronic Diarrhœa and Dysentery.

℞ Cupri sulphat.....gr. ss.
Opii pulv.....gr. ss.
Conf. rosæ, q. s. ut fiat pil.....
Take three a day. *Dr. Joy.*

In Colic.

℞ Ol. crotonis.....♏ iij.
Ol. ricini..... ℥ ij.
One tablespoonful every two hours, with opium gr. ½ to 1 three times daily. *Niemeyer.*

In Chronic Gastric Ulcer.

℞ Creasoti.....♏ iv.
Aquæ..... ℥ vj.
Misce. In tablespoonful doses. *Niemeyer.*

In Acute Gastric Catarrh.

℞ Tannin.....ʒ ss.
Aquæ destil..... ℥ iij.
Misce. A teaspoonful every two hours where great purging but no vomiting. *Niemeyer.*

In Chronic Diarrhœa and Dysentery.

℞ Cupri sulphat.....gr. vj.
Myrrhæ.....gr. xij.
Conf. rosæ.....ʒ ij.
Fiat pil. xij. Sumat unam sextis horis.
Dr. Neligan.

In Dysentery.

℞ Catechu pulv..... ʒ ij.
Pulv. acaciæ..... ℥ ss.
Aquæ..... ℥ vj.
Misce. A tablespoonful every two hours.

Niemeyer.

In Acute Nephritis.

℞ Tinct. scillæ..... ʒ ij.
Tinct. digitalis..... ʒ iij.
Tinct. ferri perchlor..... ʒ ij.
Decoct. scoparii ad..... ℥ vj.
Misce. A tablespoonful three times a day.

Dr. Grainger Stewart.

In Chronic Gastric Catarrh.

℞ Bismuthi submit.....gr. x.
Argent. nit.....gr. i—ij.
Misce. To be given at once on an empty stomach before breakfast. *Niemeyer.*

Fermentative Dyspepsia.

℞ Bismuth beta-naphthol.....gr. iij.
Salol.....gr. i.
Creosoti.....m. j.
Ol. caryophylli.....m. j.

M. Ft. tablet.

Sig.—One after meals for chronic gastro-intestinal catarrh with distress and fullness after eating.

For Indigestion.

℞ Pulv. rhei.....gr. i.
Sodii bicarb.....gr. v.
Pulv. ipecac.....gr. $\frac{1}{8}$
Ol. menth. pip.....m. $\frac{1}{20}$

M. Ft. tablet.

Sig.—One or two after meals for indigestion with sluggish liver and constipation. *Roosevelt Hospital.*

This is one of the most generally useful of anti-dyspepsia tablets, and is much more easily taken by most patients than the liquid preparations of rhubarb and soda.

For Painful Indigestion.

℞ Creosoti.....gr. $\frac{1}{8}$
Cocainæ hydrochloratis.....gr. $\frac{1}{16}$
Cerii oxalatis.....gr. ij.
Pepsini pur.....gr. $\frac{1}{4}$
Tr. nucis vomicæ.....m. $\frac{1}{4}$

M. Ft. tablet.

Sig.—One or two after meals for painful indigestion with or without flatulence.

Antifermentative.

℞ Sodii salicylatis.....gr. ij.
Pulv. zingiberis.....gr. j.
Carb. ligni.....gr. ij.

M. Ft. tablet.

Sig.—One or two after meals for *flatulent dyspepsia*, *foul breath*, *water brash*, or *chronic diarrhea.*

Colic.

℞ Spt. chloroform.....
Tr. cardamom. co.....aa ℥ ij.
M. Sig.—Teaspoonful every half hour until relieved.
Bartholow.

For Vomiting.

℞ Acid. carbol.. .....gr. iv.
Bismuth. subnitrat..... ʒ ij.
Mucil. acaciæ..... ℥ j.
Aq. menth. pip..... ℥ iij.
M. Sig.—Tablespoonful every two to four hours.
Bartholow.

For Vomiting (Non-Inflammatory).

℞ Chloroformi..... ..... ℥ ss.
Sig.—Two to five minims on sugar. *Ringer.*

For Nausea or Vomiting.

℞ Bismuthi subnitratis.....gr. iij.
Cerii oxalatis..... .....gr. iij.
Cocainæ hydrochloratis..... .....gr. $\frac{1}{12}$
M. Make into tablet or powder.
Sig.—One every half hour.

For Hæmoptysis.

℞ Acid. gallici..... ʒ ij.
Acid. sulph. aromat..... ʒ j.
Glycerinae..... ..... ℥ j.
Aq. destillat..... ..q. s. ad ℥ vj.
M. Sig.—Teaspoonful at dose; repeat frequently.
Pepper.

For Acute Diarrhea.

℞ Salol..... ʒ iss
Bismuthi subnitratis..... ʒ vj.
Mist. cretæ..... ℥ ij.
Aq cinnamomi.....ad ℥ iij.

M. Sig.—Teaspoonful every two hours following a cathartic. *Dr. F. M. Crandall.*

For Dysentery.

℞ Naphthalini.....ʒ iss.
Div. in capsules No. xviii.

M. Sig.—Two capsules every three hours.
Dr. L. E. Holt.

Diuretic.

℞ Pulv. digitalis.....gr. j.
Pulv. scillæ.....gr. j.
Potassii nitratis.....gr. ij.

M. Ft. pill.
Sig.—One every four to six hours.

The original diuretic pill so highly commended by Fothergill as to bear his name contained a grain each of digitalis, squill, and calomel. Owing to the frequency with which it caused salivation, the above formula was devised. It is a most effective diuretic. Its action may be increased by beginning treatment with small repeated doses of calomel until the bowels have acted freely. The diuretic pills may then be continued indefinitely with occasional doses of calomel.

Hemorrhoids.

℞ Acid. gallici.....gr. x.
Ex. opii.....
Ex. belladonnæ.....aa gr. iv.
Ungt. simplicis..... ℥ iv.

M. Sig.—Apply night and morning. *Dr. Hare.*

Laxative for Hemorrhoids.

℞ Potassii bitartratis .....gr. i $\frac{3}{32}$
Pulv. jalapae .....gr. i $\frac{3}{32}$
Sulphuris. .....gr. $\frac{3}{5}$
Confect. sennae.....gr. ii $\frac{3}{16}$
Potassii nitratis..... .....gr. $\frac{3}{5}$

M. Ft. pill. Sig.—One or two at night.

Basham's Mixture.

℞ Tr. ferri chlor.....f ʒ iij.
Acid. acetici dil.....f ʒ iss.
Syr. simp.....f ℥ ss.
Liq. ammom. acetat.....q. s. ad f ℥ iv.

M. Sig.—Dessertspoonful every three or four hours.
Basham.

This is a most excellent tonic for Bright's disease.

FAVORITE PERSCRIPTIONS.

For By Dr.

℞.

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

III.

DISEASES OF THE NERVOUS SYSTEM.

Neuralgia.

℞ Aconitiæ.....gr. iv.
Veratriæ.....gr. xv.
Glycerinæ ..... ʒ ij.
Cerati ..... ʒ vj.

M. Sig.—To be rubbed over the parts; care should be taken to see that there is no abrasion of the skin.

Dr. Da Costa.

Among the many local applications in neuralgia, Prof. Da Costa has found this one the best adapted for general use.

Idiopathic Epilepsy.

℞ Potassii iodidi.....
Potassii bromidi.....aa ʒ j.
Ammon. bromidi..... ʒ ss.
Potassii bicarbonat.....℈ ij.
Infus. columbæ..... ℥ vj.

Sig.—A teaspoonful before each of the three meals and three tablespoonfuls at bed-time, with a little water. *Brown-Sequard, M.D.*

When the patient's pulse is weak, substitute for the bicarbonate of potassium, in the above formula, the carbonate of ammonium, and for the six ounces of columbo an ounce and a half of the tincture of that medicine with four ounces and a half of distilled water.

Chorea.

℞ Zinci bromidi..... ʒ j.
Syr. simplicis..... ℥ j.

Sig.—Ten drops three times a day; increased as rapidly as the stomach can bear it.

Dr. William A. Hammond.

With the disappearance of the chronic symptoms the dose should be gradually diminished.

Neuralgia.

℞ Zinci valerianat..... ℈ j.
Ext. gentianæ..... ..... ℈ j.
Ext. nucis vomicæ.....gr. v.

M. Ft. pill. No. xx.

Sig.—One pill three or four times a day.

Dr. Bartholow.

In neuralgia due to reflex irritation from the female pelvic organs, this preparation of zinc is extremely beneficial.

As a Hypnotic.

℞ Chloral hydratis.....gr. xv.
Tinct. opii.....m. x.
Mist. camph..... ℥ j.

M. Sig.—Every six hours. *Fothergill.*

This is an admirable mixture to procure sleep for fever patients, without having it followed by the disagreeable effects so usual when opium is used in large doses to accomplish this purpose.

In Epilepsy in Weak and Anæmic Subjects.

℞ Potass. bromidi..... ℥ j.
Ferri bromidi.....gr. iv.
Aquæ..... ℥ ij.
Syr. simplicis..... ℥ vj.
M. Sig.—A tablespoonful twice a day.

Bartholow.

These cases are often dependent on cerebral anæmia alone, and are cured easily by using the above formula.

For Sub-Acute Mania and the Monomania of Suspicion.

℞ Hyoscyamiæ.....gr. j.
Sp. ætheris.....m. viij.
Alcohol.....m. xxiv.
Aquæ fontis.....ad. ℥ j.
M. Take at a draught. *Lawson.*

Dr. Robert Lawson, late of the West Riding Asylum, has recently made a large number of physiological and therapeutical observations on the actions and uses of the alkaloid of hyoscyamus. He has derived great benefit from the use of the above formula in recurrent, acute, and, sub-acute mania and the monomania of suspicion.

Cerebral Congestion.

℞ Sodii bromidi..... ℥ j.
Ergotæ extract. fl..... ℥ iv.
M. Sig.—A tablespoonful three times a day.

Dr. G. H. Naphey.

This formula is held by Dr. Naphey to be of great value in active cerebral congestion of all kinds, especially in the first or hyperæmic stage.

Reflex Headache.

℞ Potass. cyanid.....gr. x.—℈ j.
Aquæ lauro-cerasi..... ℥ iv.
Sig.—A compress moistened with the solution, to be applied to the seat of the pain.

Dr. Trosseau.

In reflex headache, whether gastric, cardiac, pulmonary, or menstrual, and the headache accompanying the pyretic state, from a quarter to a half hour contact with the skin of this solution usually effects a cure.

Anti-Epileptic Mixture.

℞ Sodii bromidi.....
Potass. bromidi.....
Ammon. bromidi.....aa ʒ ʒ
Potass. iodidi.....
Ammon. iodidi.....aa ʒ i½
Ammonii carbon..... ʒ i
Tinct. calumbæ.....fl. ℥ i½
Aquæ q. s. ad.....fl. ℥ ʒ
Mix. Full dose: one and a half-drachms before each meal, and three drachms at bedtime.

Dr. Brown-Séquard.

Hammond's Mixture.

℞ Quiniæ sulph.....
Ferri pyrophos.....aa ʒ i
Strychniæ.....gr. i
Acidi phos. dil. (tribas.).....fl. ʒ ii
Syrupi zingiberis.....fl. ℥ ii
Aquæ q. s. ad.....fl. ℥ iv

Mix. Dose: a teaspoonful.

Anti-Epileptic Pill.

Ninety-ninth Street Reception Hospital, New York.

℞ Argenti nitrat.....gr. x
Zinci oxidi.....gr. xx
Micæ panis.....q. s.

Mix. Divide into twenty pills. Dose: one pill thrice daily.

Nervine Tonic.

℞ Zinci valerianatis.....gr. xxiv.
Confectionis rosæ.....q. s.

Ft. mass in pil. duodecim divid. Take as required.

Farquharson.

Delirium Mixture.

℞ Potass. bromid..... ʒ iv
Tinct. valerian. amm.....fl. ℥ i
Tinct. lupuline.....
Tinct. digitalis.....aa fl. ℥ ½
Aquæ q. s. ad.....fl. ℥ iv

Mix. Dose: a tablespoonful. To be used with care and effects to be watched.

Chorea.

℞ Zinci valerianat.....ʒ ij.
Cinchoniæ sulphat.....ʒ j.
M. Ft. pil. No. xx. Sig.—One pill thrice daily.

Dr. Da Costa.

Cannabis Mixture.

Insane Asylum, Blackwell's Island, N. Y.

℞ Tinct. cannabis ind.....m. 10
Spiritus menthæ pip.....m. 1
Aquæ q. s. ad.....fl. ʒ 1
Mix. One dose. To be taken thrice daily after meals.

Tincture of Phosphorus.

Bellevue Hospital, New York.

℞ Phosphori.....gr. 32
Alcoholis absol.....fl. ℥ 46
Tinct. vanillæ.....fl. ℥ 1
Ol. aurantii cort.....fl. ʒ 3
Alcoholis absol. q. s. ad.....fl. ℥ 48

The phosphorus is digested with the absolute alcohol, with the exclusion of air, until dissolved; then the flavoring ingredients are added, and finally the bulk is made up with absolute alcohol to 48 fl.oz.

12 fl. drachms contain 1 grain of phosphorus.

30 minims contain $\frac{1}{24}$ grain of phosphorus.

Dose: 20-40 minims, corresponding to $\frac{1}{36}$-$\frac{1}{18}$ gr. of phosphorus.

Epileptic Belladonna Mixture.

Insane Asylum, Blackwell's Island, New York.

℞ Potass. bromidi ..... gr. 25
Tinct. belladonnæ ..... m. 5
Aquæ q. s. ad ..... fl. ʒ i

Mix. One dose. To be taken thrice daily.

Phosphorus Mixture.

℞ Phosphori ..... gr. i
Alcoholis absoluti ..... fl. ℥ ʒ 5
Glycerinæ ..... fl. ℥ ʒ 1½
Alcoholis ..... fl. ʒ 2
Spts. menthæ pip ..... fl. ʒ i

Dissolve the phosphorus in the absolute alcohol by the aid of a gentle heat; then add to it the glycerine, alcohol and spirits of peppermint, previously mixed and slightly warmed.

One fl.oz. contains $\frac{1}{20}$ gr. of phosphorus.

Dr. J. Ashburton Thompson.

Compound Syrup of the Hypophosphites.

℞ Calcii hypophosphitis ..... gr. 256
Sodii hypophosphitis ..... gr. 192
Potassii hypophosphitis ..... gr. 128
Ferri sulphatis ..... gr. 185
Acid hypophosphorosi (1.036) ..... fl. ʒ 9
Sacchari ..... ℥ ʒ 12
Aquæ q. s. ad ..... fl. ℥ ʒ 18

Mix. Dose: a teaspoonful.

Sedative Mixture.

Insane Asylum, Blackwell's Island, New York.

℞ Chloralis.....gr. 15
Extr. conii sem. fl.....
Extr. hyoscyami fl.....aa m. 15
Aquæ q. s. ad.....fl. ʒ i

Mix. One dose. To be taken thrice daily, after meals.

Epileptic (Conium) Mixture.

Insane Asylum, Blackwell's Island, N. Y.

℞ Potass. bromidi.....ʒ ½
Ext. conii fl.....m. 15
Aquæ q. s. ad.....fl. ʒ i

Mix. One dose ; to be taken thrice daily.

Gross's Neuralgic Pills.

℞ Quiniæ sulphat.....ʒ ii
Morphiæ sulphat.....gr. 3
Strychniæ.....gr. 2
Acidi arseniosi.....gr. 3
Extr. aconiti.....gr. 30

Mix. Divide into 60 pills.

In Neuralgia.

℞ Tinct. cannabis indicæ.....ʒ j.
Mucilag. acaciæ.....℥ j.
Syr. zingiberis.....℥ ss.
Aquæ menth. piperitæ.....ad. ℥ vj.
Ft. mistura et.....

Sig.—Two tablespoonfuls every four hours.

Farquharson.

Brown-Sequard's Neuralgic Pills.

℞ Extract hyoscyami.....
Extract conii.....aa gr. 40
Extract ignat. amar.....
Extract opii.....aa gr. 30
Extract aconiti.....gr. 20
Extract cannab. ind.....gr. 15
Extract stramonii.....gr. 12
Extract belladonnæ.....gr. 10
Mix. Divide into 60 pills.

Narcotic.

℞ Ext. belladonnæ.....gr. iij.
Camphoræ.....gr. xij.
Ext. hyoscyami.....gr. xv.
M. Make six pills, of which take one when going to bed. *Farquharson.*

Hypnotic.

℞ Potass. bromid.....gr. xxx.
Syrupi aurantii..... ʒ j.
Aquæ aurantii flor..... ℥ ij.
To be taken at a draught at the time sleep is desired. *Farquharson.*

Epileptic (Ergot) Mixture.

Insane Asylum, Blackwell's Island, N. Y.

℞ Potass. bromidi.....
Ammon. bromidi.....aa ʒ ½
Ext. ergotæ fl.....m. 15

Mix. One dose: to be taken thrice daily, in cases characterized by considerable maniacal excitement following the attack, indication of cerebral congestion, and especially where hemorrhage is feared.

Dr. Chas. R. Smith.

To Allay Craving for Drink in Dipsomaniacs.

℞ Tinct. capsici.....m. x.
Tinct. nucis vomicæ.....m. x.
Acid. nit. dil.....m. xx.
Aquæ.....ad. ℥ j.

To be taken at a draught three times a day.

Farquharson.

Tonic in Neuralgia.

℞ Ammon. chlorid..... ʒ ii.
Ext. glycyrrhizæ..... ℥ ss.
Syr. tolut..... ℥ j.
Aq. cinnamoni.....ad. ℥ viij.

Sig.— ℥ j. every four hours. *Dr. Farquharson.*

Epilepsy.

The following prescription is a useful one for a child about fourteen or fifteen years:

℞ Pot. bromid..... ℥ ss.
Ferri bromid.....gr. ij.
Glycerini ..... ℥ iv.
Aquæ.....ad. ℥ viij.

Mix. A tablespoonful twice a day. *E. Ellis.*

In Hysteria.

℞ Auro-sodii chloridi.....gr. v.
Pulv. tragacanth.....ʒ j.
Sacchar. alb.....q. s.
Misce, fiat pil. xl.

One pill an hour after dinner and supper. After, two pills following these meals, and gradually increase to eight pills daily. *Niemeyer.*

In Insomnia.

℞ Potass. bromid.....ʒ vj.
Aquæ destil.....℥ v.
Misce. Three teaspoonfuls before dinner, and four at bedtime, with a little wine and water.

Dr. Brown-Séquard

Idiopathic Epilepsy.

℞ Ext. ergot. fl.....½ to 1 drachm.
Ext. conii fl.....5 drops.
M.—Sig. Three times daily.

Dr. E. C. Spitzka, New York.

Infantile Convulsions.

First give a purgative dose of calomel, and follow in a few hours by

℞ Chloral hydrat.....4 grains.
Potassii bromidi.....8 grains.
Aquæ.
Syrupi, of each.....1 drachm.
M.—Sig. Dose for a child two years old.

Dr. A. Jacobi, New York.

HEADACHE.

Treatment.—(1) *Of Sick-headache.*—Looking to the true cause of sick-headache as one which lies deep in the patient's idiosyncrasy, and as an affection which is developed by a hundred different sources of excitement, the advice to sufferers is to give as much tone as they can to their nerves by adopting all those methods which experience has shown to be good, and then avoid, as far as is practicable, all those causes which are known to excite an attack. Purgation has no good result. The only remedies which are of any avail are those which act on the nervous system, such as hot tea and coffee; or, after the stomach is quieter, and the more urgent symptoms have passed off, a little wine or ammonia. If the headache take more the form of hemicrania, then other remedies are occasionally useful, as the local application of the bisulphide of carbon, or galvanism, and internally the bromide of potassium. This is the only drug which Dr. Wilks has really seen to be serviceable. Whilst the nausea exists and the worst symptoms prevail, even this remedy is of no avail. As regards tea and coffee, which often relieve, it is possible that these and other stimulants, taken in excess, render the nervous system more susceptible to the attacks.

In the treatment of the paroxysm nothing is so ef-

ficacious as complete rest in the recumbent posture, with warmth, in a perfectly quiet and darkened room. A cup of very strong tea or coffee has also given relief. A comparatively new drug—guarana—has also been recommended. A full dose of brandy, or of ammonia, has also sometimes dispelled the headache; so has continued sipping of a solution of carbonate of potash or of soda. Pressure upon the carotid artery of the affected side sometimes also relieves the pain, but for a time only. Evaporating lotions, belladonna, or aconite ointment may also ward off or subdue the headache; but, as a rule, time is the most efficient element, with warmth to the feet and to the body, and the induction of sleep.

Adopting the theory of Dr. Latham, what are the indications for treatment with which it supplies us—(*a.*) During the stage of disturbed sensation; (*b.*) during the stage of headache; and (*c.*) during the intervals between the attacks? The following are the results of his experience:

(*a.*) *During the Stage of Disturbed Sensation.*—It is found that the longer this stage lasts, the greater will be the headache; and therefore it is desirable to shorten it as much as possible. If the condition depend upon deficient supply of blood to a part, such means must be adopted as shall assist and increase the flow of blood to the part; and this can be done in some measure by posture and stimulants. Directly the glimmering appears, the patient should lie down with the head as low as possible, and if the glimmering be on the right or left of the field of vision, he should lie on the opposite side. Let him take at once one

ounce of brandy, either "neat" or in half a bottle of soda-water. Champagne would be preferable, being more diffusible; but its administration would often involve a little delay. If alcoholic stimulants be objected to, or if it be not advisable to recommend them, then a teaspoonful of sal-volatile in water may be prescribed instead. If the patient be chilly or his feet cold, the couch should be drawn before the fire, a hot bottle applied to the feet, and his body covered over with a warm blanket or eider-down quilt. By these means the heart is enabled to drive the blood with greater force to the brain, and the duration of the vibratory movement is thereby materially lessened. After it has passed off, the patient should lie still for a time, so that the glimmering may not return. This injunction will only be necessary when the headache is slight; if it be severe, attended with much nausea or vomiting, the patient will be little disposed or able to leave the recumbent position. If, instead of the disturbance of vision preceding the headache, there be a feeling depression or irritability (fidgets), the administration of such cerebro-spinal stimulants as henbane, valerian, assafœtida, spirits of chloroform, or ether, will often cut short the attack; ten or fifteen drops of the tincture of henbane, with the same quantity of spirit of chloroform, will soothe the nervous irritability in the slighter forms, and may be repeated in three or four hours, if necessary. If there be great mental depression, then valerian or assafœtida should be tried. Stillé says,—"Nothing is more astonishing in the operation of remedies than the promptness and certainty with which a dose of valerian or assafœtida

dispels the gloomy visions of the hypochondriac, calms the hurry and agitation of nervous excitement, allays commencing spasms, and diffuses a soothing calm over the whole being of one who but an hour before was a prey to a thousand morbid sensations and thick-coming fancies of danger, wrong, or loss." Dr. Latham gives the preference to valerian, and prescribes from half a drachm to a drachm of the ammoniated tincture. Or it may be combined, as in the following formula, with bromide of soda; ammoniated tincture of valerian ℥ iv. with ʒ vi. bromide of sodium, in teaspoonful doses every hour. The assafœtida may be given in the form of the spiritus ammoniæ fœtidus of the Pharmacopœia, also in half drachm or drachm doses. As a rule, alcoholic stimulants are not advisable in this stage of the headache. A small quantity will cause flushing, heaviness, slight confusion of thought, etc., without relieving the depression; and though the severe headache may be averted, alcoholic stimulants do not answer so well as the remedies previously mentioned.

(*b.*) *During the Stage of Headache.*—If the headache be slight and the patient soon able to sit up, there is little to be done; a cup of coffee or tea, cheerful conversation, a walk, drive, or ride, may often help to remove the pain. If, however, the headache, and nausea be severe, then the administration of further remedies is called for. The patient should keep perfectly still and quiet, with the room darkened; for every sound or sight causes pain, and the slightest movement is sufficient to produce gastric uneasiness. Sometimes free evacuation of the contents of the stomach, espe-

cially if it contain undigested food, is followed by relief. Dr. Fothergill says,—"An emetic and some warm water soon wash off the offending matter, and remove these disorders," which may be very well where there is any offending matter to wash off, but it is not very often that this is the case; the nausea frequently continues long after the contents of the stomach have been discharged; an inverted action of the duodenum is set up; the bile appears in the fluids excreted; the patient believes that all his troubles are due to "its overflow;" "it's all liver," he says, and it is sometimes difficult to persuade him to the contrary. Generally, then, you should try to relieve and check the vomiting. Iced soda-water, with or without two or three drops of dilute hydrocyanic acid, or spirit of chloroform; cold tea; the effervescing citrate of potash, with hydrocyanic acid, may often afford marked relief. The headache may be lessened by applying cloths dipped in cold water, or evaporating lotions, to the head; if the extremities be cold and the headache severe, a warm stimulating foot-bath can be tried so soon as the nausea will allow the patient to sit up. If the attacks occur in the early part of the day, as soon as the pain has subsided it is generally better for the patient to sit up, or move about, or take exercise in the open air. During the attack the appetite is diminished, the idea even of food provoking disgust. Still, after the nausea has passed away and the headache has continued a few hours, a plate of hot soup or some easily digested food will often have a good effect in equalizing the cerebral circulation. A remedy which may very often be given with advan-

tage if the headache be severe, is bromide of potassium, in doses of five, ten, or fifteen grains, to which thirty or forty minims of sal-volatile may, in some cases, be added with advantage; and if the nausea still continue, these may be given in combination with the effervescing citrate of potash. A saline purgative at the commencement of an attack is sometimes an effectual remedy; but, as a rule, the use of purgatives is objectionable. So far, these measures are only palliative.

(*c.*) *During the Intervals between the Attacks.*—First of all, the exciting cause is to be discovered and removed. Hours of study or work must be abridged, excessive bodily fatigue, loss of rest, everything, in fact, must be avoided which the sufferers know from individual experience will act as exciting causes. Where the attacks are associated with excessive mental work, they should be regarded as danger-signals, showing necessity for relaxation. The tone of the bodily and nervous systems must be improved by proper medicinal and hygienic means; and the chief remedies are steel, strychnine, phosphorus, and cod-liver oil. The success, however, following these remedies depends a great deal upon the way in which they are administered. For a day or two after the attack the stomach and bowels may possibly be disordered, and not in a fit state to tolerate such remedies. This must first be corrected. The simple vegetable bitters, such as gentian, with small doses of henbane and some aromatic, may be of service; and, if necessary, one or two grains of blue pill, with four or five of compound rhubarb pill, may be given at night. We may

then try steel. In anæmic cases it may be advisable to stimulate the action of the heart, as by the following:

℞. Ammoniæ muriatis, ℥ ss., Tinct. acteæ racemosæ, aquæ, a a ℥ iii. Two fluid drachms for a dose after meals in a wineglass of water.

If there be despondency with depression of spirits, phosphorus to the extent of $\frac{1}{100}$ to $\frac{1}{50}$ of a grain, combined with nux vomica or carbonate of iron, or both taken with each meal, will improve the tone of the patient. If the attacks have been frequent, or if there be any scrofulous tendency, the iodide of iron may be given in the following form:

℞. Ferri et ammon. citrat., gr. v.; potassii iododi, gr. ij.; aquæ, ℥ j.; and, according to circumstances, fifteen to twenty minims of tincture of henbane or twenty or thirty minims of aromatic spirit of ammonia may be added.

If the stomach be at all irritable, this medicine may be given in the effervescing form, adding to each dose twenty grains of bicarbonate of potash, and directing it to be taken with a tablespoonful of lemon-juice, or a corresponding amount of citric acid: the dose to be taken twice a day, about eleven and four. The effervescing form may soon be left off, and then to each dose may be added five minims of liquor of strychniæ (omitting the henbane and sal-volatile, and continuing the iodide of potassium according as it seems to be indicated or not). In other cases, the citrate of iron and ammonia with strychnine may be given at the beginning, sometimes combined with infusion of calumba. The iron is indicated by the

greater or less anæmia of the patient; but the strychnine is a very important remedial agent in the disorder. In small doses it acts as a simple tonic, increasing the appetite and improving the digestion ; it dilates the vessels and, thus increasing the supply of blood, it augments the activity of the spinal cord (Harley). It promotes the capillary circulation, and therefore its use is advisable for persons troubled with cold hands and feet (Anstie); and if it fulfil these conditions, it is clearly indicated in this disorder. Cod-liver oil also often acts very beneficially. "It has been found by experiment that great exertion and prolonged labor can be endured without fatigue when starchy and fatty foods are alone eaten; . . . . and there is reason to think that cod-liver oil is more easily absorbed than other similar substances" (Ringer). "It improves the digestive process, increases the proportion of red corpuscles in the blood, and invigorates the whole nutritive function" (Wood); and it particularly sustains the energy of the brain during prolonged mental exertion. A gentleman in the foremost rank at the bar told Dr. Latham that whenever he was engaged in a jury-trial which was likely to tax his energies to a greater degree than usual, the thing which best sustained him was a good dose of cod-liver oil taken in the morning before going into court ; and others engaged in mental work have confirmed this view. Besides its other properties cod-liver oil is a nutrient and tonic in its action on the cerebro-spinal nervous system. As a remedy for these nervous headaches, Dr. Latham prescribes it once a day, beginning with a small teaspoonful imme-

diately after breakfast, and gradually increasing the quantity to a tablespoonful, but not beyond, unless in exceptional cases.

The action of the bowels must be regulated, but not by strong purgatives. Five grains of the Socotrine aloes pill, given at night, are generally sufficient. If the bowels be habitually constipated, then no remedy seems to answer so well as the aloes and iron pill. Five grains given twice a day, half an hour before meals, will act freely; and in a few days the dose will have to be diminished, for its effeet is augmented instead of being lessened by continual administration, especially when strychnine is given at the same time. The natural waters of Friedrichshall or Marienbad may, in many instances, be of service, given as laxatives.

In 1872 Dr. Samuel Wilks, Physician to Guy's Hospital, directed the attention of the profession to guarana as a remedy for sick-headache, and at the same time asked for the experience of those who already had some acquaintance with the drug. His own knowledge of it dated about 1870, when, after the appearance of a lecture of his upon sick-headache, he received a letter from Mr. Helmcken, of British Columbia, in closing two powders, which he recommended with much confidence as able to cure the complaint. "Upon my first headache after the receipt of Mr. Helmcken's letter, I took the powder, but with only doubtful effect. I therefore did no more than casually mention the medicine to my friends, but did not recommend it. A few weeks ago, after the appearance of a second communication of mine in the jour-

nal upon the same complaint, I received a letter from Dr. Wood, of Montreal, in which he also recommended guarana as a remedy for headache, and gave a history of his own personal sufferings and the relief which he obtained. He says, 'By taking one of these powders and remaining quiet when I have felt premonitory symptoms by a beginning of pain always in the right temple (headache on the other side, or in any other part of the head, I never mind), I have carried off the attack; and, with the first box, absolutely put it off for two months—something which had never occurred in my life before.' Upon so good authority, I determined to try the remedy in a more systematic manner, and requested my neighbor, Mr. Hooper, the chemist, to procure me a packet of the powders. These I have recommended to several patients and friends; and the result is so encouraging, that I have hastened to suggest their trial to my professional brethren. One lady speaks most enthusiastically of their power, as she has now, on two separate occasions, had her headache arrested by their use. The drug has long been known, for mention is made of it in English and French pharmacologies, but appears never to have come into general use. It consists of the seeds of a tree growing in Brazil, called Paullinia sorbilis; and these, according to Johnstone, in his "Chemistry of Common Life," are used as we do cocoa. The seeds are ground into powder, and contain an alkaloid which is said to be identical with that found in tea and coffee. The medicine is manufactured by Grimault & Co., No. 7 Rue de la Feuillade, Paris" (Wilks, *Brit. Med. Journ.*, April 20, 1872).

The general outcome of experience regarding the treatment of sick-headache seems to be that—

(1.) Bromide of potassium, Indian hemp, Guarana or Paullinia powder, are remedies which have afforded relief. With reference to bromide of potassium, it is so valuable a medicine in many cases of sick-headache that it can scarcely be superseded by a better remedy. The patient may have a splitting headache, fatigued and worried after a hard day's work. After fifteen or twenty grains of bromide of potassium, he presently goes off to sleep in his easy chair, and wakes in an hour well (Wilks). It is a medicine which Dr. Wilks always employs first, having seen such eminent advantages follow its use. "I have known (he writes) many patients declare, that the bromide was the first medicine they had taken in their lives which had the slightest effect in relieving their headache." As to Cannabis Indica, it seems best adapted to the interval between the attacks, in doses of a few drops of the tincture, when headaches are recurrent. It is best given in ten-minim doses, three times a day, continued for weeks or months (Wilks, John Murray). As to guarana powder, it seems most of all useful with females in arresting headache, so that it must be taken early when the headache is approaching; and if headache is not arrested, a second dose should be taken in fifteen, twenty, or thirty minutes after the first. If the medicine is of use, the headache will have gone in about an hour. The dose is 15 grains of the powder; and as it is somewhat astringent, it might be combined as an electuary with some laxative, such as the senna electuary (Hollis).

(2.) Certain classes of cases ought to be distinguished from each other. There are two principal types: the congestive, met with in dark or plethoric people, in those with constipated bowels and suppressed secretions, and those addicted to alcoholic beverages, or leading a monotonous sedentary life; and that arising from exhaustion, whether from want of sleep or want of food, or great bodily or mental fatigue, strong emotions, etc.,—this may be nervous, or asthenic, or anæmic in many cases. This latter form is very common in slight, delicate, and tuberculous females, and sometimes in the male sex. The treatment of these two forms must, of course, differ widely. An emetic or a purge in the one case will do what only rest, food, stimulants, and such medicines as quinine, iron, opium, hyoscyamus and belladonna may afford in the other case (Dr. Bathurst Woodman). Rest and quiet, especially sleep, and abstinence from food, give relief and cure by a natural process of restoration. Generally, no relief is obtained till after a night of sound sleep—*i. e.*, "one night through the sheets"—awakening next morning free from pain, but generally more or less depressed. If depression continue, phosphorus with nux vomica in the form of sugar-coated or pearl-coated pills may be given to the extent of one fiftieth of a grain of phosphorus and one eighth of a grain of extract of nux vomica three times a day with food. Such despondency and depression are apt to occur in those who are over-worked mentally; or who are harassed by business; or who suffer great mental anxiety. If there be sleeplessness, the following pill may be of use:

℞ Camphoræ pulveris, gr. xx.; Ext. Cannabis Indica, gr. xii.; Ext. Hyoscyam., gr. xxiv.; misce bene. Divide in pill xii. signa, one at night, to be repeated in two hours, if necessary, to procure sleep.

As to local application, a large sinapism over the stomach, followed by a warm linseed meal poultice, has sometimes been of service. If pressure relieves, the head may be bound round with a handkerchief. When the head throbs and is hot, instinct seems to suggest cold and pressure to the part; and one of the best external appliances on which we can rely to procure relief is the wet bandage tied tightly round the head. The method must be instinctive, for it is universal, and has been used from all time. Shakespeare is often quoted to illustrate the morbid states of the body as well as the passions of the mind, and he also testifies to this ancient practice. For example, in the scene between Hubert and Arthur in "King John," the latter, when petitioning for the preservation of his eyes, says:

> "When your head did but ache
> I knit my handkerchief about your brows."

And in Othello we have not only the remedy for headache given, but the cause. The former was the handkerchief about which the chief interest of the play centred.

> "*Desdemona*, Why do you speak so faintly?
> Are you not well?
> *Othello*, I have a pain upon my forehead here.
> *Desdemona*, Faith, that's with watching: 'twill away again.
> Let me but bind it hard, within this hour
> It will be well."

Sick-headache is not to be cured by gastro-hepatic remedies. It is a purely nervous affection, due to hereditary predisposition, and excited by causes innumerable which act on a susceptible nervous system. There is, then, no cure in the proper sense of the term, for this would imply a change in the patient's nature; and for the attacks themselves, when severe, the only relief which can be reckoned upon is to be found in a wet bandage round the head, profound quiet, and, if possible, sleep (Dr. Bathurst Woodman).

Considerable relief may be obtained from the use of cold applications, in the form of wet rags, ice in a bladder, or a little of the ether-spray; but in not a few cases, after all, the best treatment during the attack is absolute rest and quiet, in a dark room, with warmth to feet and body, total abstinence, except perhaps from fluids (hot and alkaline) in small quantities at a time; to go to bed at once, and refrain from the use of any remedies.

In a lecture on the therapeutics of headache, delivered at Bellevue Hospital Medical College, Dr. A. A. Smith suggests certain combinations of remedies which he has found of service in certain forms of headache: (1.) Malarial headache, where the pain is most frequently located in the sub-occipital or frontal regions, or on either side,—hemicrania. If distinctly periodical, give ten or fifteen grains of quinine two or three hours before the expected attack. It may be necessary to push the quinine in divided doses until cinchonism is produced, and kept up for several days. Then gradually diminish the dose. If the pain still continues to recur, resort to arsenic and belladonna,

five-drop doses each of Fowler's solution and belladonna tincture, after meals, increasing the Fowler's one drop each day, until œdema arsenicalis is produced. This will seldom fail to give relief.

(2.) In Headache dependent on Gout:

℞ Vini colchici seminum, ʒ iij.; Lithii bromidi, ℥ ss.; Syr. zingiberis, ℥ ss.; Aq. cinnamonii, q. s. ad. ℥ vj.; M. Sig. ℥ ss. in a tumbler of Vichy water every four hours.

(3.) Rheumatic Headache is characterized usually by tenderness of the scalp, which is increased on pressure or motion. Use the mild Faradic current on the scalp, and internally the following:

℞ Potassii iodidi; Ammoniæ muriatis, aa ʒ jss.; Infusi humuli, ℥ vj.; M. Sig. ℥ ss. four times a day in a wineglass of water.

In cases which have not yielded to the above treatment, bromide of ammonium in twenty-grain doses every two hours may be effectual.

(4.) Uræmic headache is of great importance as a symptom of serious disease. The pain in the head may be the first evidence that there exists renal disease, and that we have to deal with uræmic headache. The judicious plan of treatment in such cases is to call into action one or all of the three great emunctories of the body, the kidneys, the intestines, and the skin. To make the kidneys act, apply dry cups over the region of them, and give internally the following:

℞ Potassæ acetatis, ʒ vj.; Infusi digitalis, ℥ vj.; M. Sig. ℥ ss. every third hour.

The infusion should be made from fresh English

leaves. Give this until the kidneys act freely, if you can make them do it within twenty-four hours. If the kidneys do not act freely, and the headache be not relieved within twenty-four hours, give a saline cathartic. A treatment almost domestic, and often very effectual, is to put an ounce of cream of tartar in a quart of water, and have the patient drink this in eight or ten hours. It acts both as a diuretic and cathartic. Do not use hydrogogue cathartics unless convulsions be threatened. Some prefer elimination by the skin. This can be done by diaphoretics and the hot, moist, or dry air baths. Diuretics, cathartics, and diaphoretics are only to be used where there is deficient quantity of urinary secretion. There are other causes of headache in Bright's disease which occur independently of the presence of an abnormal amount of urea in the blood, and which yet are dependent on the results of the kidney disease. These causes may be anæmia, neuralgia, œdema of the brain itself, serous effusion into the ventricles; and, in acute Bright's disease, cerebral congestion. Under the last condition, if the headache be very severe and convulsions threaten, blood may be taken if the patient's condition will admit of it, from twelve to twenty ounces, if necessary to relieve distressing symptoms. The best way to take it is by means of wet cups over the region of the kidneys. If the headache be dependent on serous effusion into the ventricles, or on cerebral œdema, improve the vitiated condition of the blood, and stimulate the heart and kidneys by acetate of potash and infusion of digitalis. There is apt to be general anasarca with this effusion and œdema.

(5.) The Headache of Acute Alcoholism, or inebriety. The first indication is to remove the alcohol from the intestinal canal. For this, give of rhubarb and magnesia calcined each a half drachm, and then the following:

℞ Spiritus ammoniæ aromat., ʒ ij.; Tincture camph., ʒ jss.; Tincture hyoscyami, ʒ ijss.; Spiritus lavandulæ comp., q. s. ad ℥ ij.; M. Sig. ʒ j. every hour until the headache is relieved, and then give capsicum gr. ij. and quinine gr. iij. before each meal for several days. If there be sleeplessness give ℞ Sodii bromidi, ℥ ss.; Chloral. hydrat., ʒ ijss.; Syrupi aurantii cort., ℥ ss.; Aquæ, ℥ iijss. M. Sig. ℥ ss. at night; repeat in two hours if necessary to produce sleep.

(6.) *Dyspeptic Headache.*—If there be indigestible food in the stomach, and it have been there some time, give an emetic, as mustard and warm water, or sulphate of zinc, gr. xv. If there is evidence of indigestible food in the alimentary canal beyond the stomach, give gr. xx. of rhubarb and magnesia each, to remove it from the bowels. If the headache be frontal, and the pain be located immediately over the eyes, give dilute nitro-muriatic acid in ten-drop doses, well diluted after meals. If the pain be located about the roots of the hair, give an alkali before meals, as gr. xv. of bicarbonate of soda or magnesia. The dyspeptic headache oftentimes is not confined to these regions, but spreads over the entire head. In such cases combine an acid with an alkali, and add to these nux vomica, as in the following prescription:

℞. Sodæ bicarbonatis, ʒ ijss.; Acidi nitro-mur. di-

luti., ʒ ij.; Tinct. nucis vomicæ, ʒ jss.; Syr. aurant. cort., ʒ vj.; Aquæ, q. s. ad ℥ vj. M. Sig. ℥ ss. after meals in a wineglass of water.

If there be gastric pain, a mild counter-irritant, âs a mustard plaster to the epigastrium, will often relieve the pain in the head as well as the pain in the stomach. If flatulence be a troublesome symptom, give the following:

℞. Bismuthi subcarbonatis, ʒ ijss.; Tinct. nucis vomicæ, ʒ jss.; Tinct. cardamomi comp.; Spiritus lavandulæ comp. aa q. s. ad ʒ iv. M. Sig. ʒ ij. before meals in a wineglass of water.

If there be constipation, the following pill may be given, in the morning:

℞. Aloes pulv., ʒ ss.; Ext. nucis vomicæ, gr. v.; Ext. belladonnæ, gr. iv.; M. Div. in pilul. xv.

In some forms of headache associated with stomach indigestion, small doses, often repeated, of tincture of nux vomica may be effectual. A single drop every fifteen minutes and continue this two or three hours if necessary. In other cases, where the headache comes on soon after a meal, and seems to depend on delayed stomach digestion, large doses of pepsin are effectual. Give a half drachm of saccharated pepsin in a wineglass of sherry wine, three times a day, and let it be taken during meals.

(7.) *Congestive Headache.*—Cerebral congestion as a cause of headache may be divided into two varieties, active and passive. These demand almost directly opposite plans of treatment. In the active variety the patient should be kept in a darkened room, perfectly quiet, and have cold and evaporating lotions applied

L. of C.

to the head. A saline cathartic may be given; and the following prescription:

℞. Sodii bromidi, ʒ ijss.; Fl. ext. ergot., ʒ ijss.; Syr. zingib., ℥ ss.; Aq. aurant. flor. q. s. ad ℥ iv. M. Sig. ℥ ss. every two hours.

If the skin be hot and dry, and the pulse full and rapid, give two drops of Fleming's tincture of aconite root every two hours, until the heart's action is sensibly diminished. Sometimes also a hot mustard foot-bath will give relief. The passive congestive variety demands a different mode of treatment. It is often found associated with cardiac disease, and most frequently where there is predominant dilatation. Hypertrophy gives rise to the active variety. Improve the condition of the blood by the use of iron, quinine, bitter tonics, alcoholic stimulants, good food, and stimulate the heart's action by the use of the following:

℞. Tinct. digitalis, ʒ iij.; Spirit. amm. aromat., ʒ vj.; Spirit. lavandulæ comp.; Syr. simp. a a q. s. ad ℥ iij. M. Sig. ʒ j. every four hours.

(8.) *Anæmic Headache.*—Cerebral anæmia produces a headache which is often mistaken for the passive cerebral congestive form. It is associated with general anæmia, nervous exhaustion, and may occur in heart disease, in consequence of enfeebled heart power, such as is met with in enlargement with dilatation, fatty degeneration, and myocarditis. Improve the general condition of the patient, and stimulate the heart's action as recommended in the passive cerebral congestive variety. Nitrite of amyl will relieve the immediate headache. Let the patient inhale three to five drops of it on a piece of cotton, placed within one

nostril, while the other is held closed. When associated with nervous exhaustion, employ the following remedies:

℞. Strych. sulph. gr., ss.; Tinct. ferri chloridi, ʒ ij.; Glycerinæ ℥ ss.; Infusi gentian., q. s. ad ℥ vj. M. Sig. ℥ ss. after meals, in a wineglass of water.

Alcoholic stimulants are beneficial in headache dependent on cerebral anæmia. Champagne as a remedy, and not as a beverage, may be given to those who suffer from nervous exhaustion; or a tablespoonful of brandy, diluted with water, after each meal.

(9.) *Cerebral Tumors* give rise to headache, often severe. In all such cases, give iodide of potassium; for it cannot be safely said that in any given case the tumor does not depend on syphilis, and by administering the remedy the patient gets the benefit of the doubt.

There is reason to believe, too, that patients with cerebral tumors, dependent on other and unknown causes, are benefited by the use of iodide of potassium. If there be much sleeplessness, use the bromide and chloral mixture. Ergot has also been used. It is usually given in large doses, beginning with ʒ i. of the fluid extract three times a day, and gradually increasing the dose to half an ounce. *W. Aitken.*

But little can be added to this admirable article as regards the preventive and curative treatment of headaches. Since it was written, however, there have been introduced several drugs effective in affording temporary relief. There is, it must be said, a strong tendency at the present day to rely too much upon these palliative measures to the neglect of the more permanent treatment ad-

vocated by Aitken. However, several of these are very valuable aids in the treatment of some of the most stubborn and discouraging conditions the practitioner is obliged to meet.

Antipyrin was the first of these drugs to come into general notice. It is an efficient remedy in migraine and other forms of headache, but owing to uncertainty of action and a tendency to depress the heart and cause urticaria it is not now extensively used. In the hands of the laity, it has undoubtedly caused much harm. The following is a very effective prescription for migraine:

℞ Caffeinæ citrat..... .....gr. xl.
Sodii bromid..... ʒ iv.
Antipyrin..... ʒ ij.
M. Div. in chart. No. xx.
Sig.—One powder in water as needed. *Hare.*

Acetanilid has stood the test of ten or twelve years and its place in therapeutics is well established. Bartholow sums up the experience of most practitioners in the statement that its special field of utility is the relief of inflammatory conditions of the nerves. In neuritis he has found it so effective that it has seemed to have specific action. The pains of neuralgia, myalgia, etc., due to inflammation of the nerves, are as a rule soon relieved by doses not exceeding three to five grains three or four times a day. It is one of the few drugs which have any controlling influence on the pains of locomotor ataxia. In migraine it is particularly effective, as well as in the neuralgias of malaria. An effective and popular combination

is known as "migraine tablets," consisting of acetanilid, camphor monobrom., and caffeine. This combination is made up in numerous strengths, one of the most generally useful being the following, known as Migraine No. 1:

℞ Acetanilid.....gr. ij.
Camph. monobrom.....gr. ss.
Caffeine citrat.....gr. ss.

M. Ft. tablet.

Sig.—One, repeated every hour for four or five doses for *migraine*, *neuralgia*, and *sick headache*.

Phenacetin is used in most of the conditions described under acetanilid. The dose is a little larger than that usually given of acetanilid; ten grains is equal to fifteen grains of antipyrin. It is believed by many to be more effective in relieving pain than acetanilid and to be less depressing. While this is doubtful, it certainly does not cause cyanosis so readily; its great expense is perhaps the chief reason for its being less used than acetanilid.

Lactophenin is effective in some forms of rheumatic headaches, but for most forms of headache is less effective than the preceding three drugs.

For Locomotor Ataxia.

℞ Antipyrin.....ʒ j.
Syr. zingiber.....ʒ j.
Aquæ.....ad ℥ iv.

M. Sig.—A teaspoonful every one to four hours for three to six doses for the lightening pains.

Germain Sée.

Nervine Tablets.

℞ Pulv. camph.....gr. j.
Ext. hyoscyomi.....
Ext. valerianæ.....aa gr. ss.

M. Ft. tablet (chocolate-coated).

Sig.—One every half hour for three doses, then every two hours

An efficient combination for the treatment of *neurasthenia*, *hysteria*, *insomnia*, and *headache*, particularly when associated with uterine or ovarian disease. It is also efficacious in *ovarian neuralgia*. Some practitioners add one-half grain of sumbul to each tablet.

For Hysteria.

℞ Ext. sumbul.....
Ferri sulphat. exsic.....aa gr. xx.
Pulv. asafœtidæ .....gr. xl.
Acid. arsenosi.....gr. ss.

M. Ft. pilulæ No. xx.

Sig.—One or two pills thrice daily. *Goodell.*

This is particularly efficacious for hysterical, anæmic women, suffering from ovarian and uterine disease.

For Acute Mania.

℞ Paraldehyde.....f℥ss.

Sig.—Thirty to fifty minims in water by the rectum.

Ringer.

For Epilepsy.

℞ Potassi bromidi.....℥ j.
Sodii bromid.....℥ ss.
Ammonii bromid.....ʒ ij.
Syrup.....f℥ ij.
Aq. Gaultheriæ.....q. s. ad f℥ vj.

Sig.—A teaspoonful three times a day for a child of seven years. *Rex.*

Insomnia.

℞ Sulphonal.....gr. xxx.
Syrupi.....℥ ij.
Mucilag. acaciæ.....ʒ ij.
Aquæ.....q. s. ad ℥ j.

M. Sig.—Half to all of this at one dose, as may be required. *Hare.*

OR

℞ Trional.....gr. xv.-xx.

Sig.—For one dose at bed-time.

Trional acts much more promptly than sulphonal, and usually causes less drowsiness during the following day. It is especially indicated in case of extreme nervousness. It has little or no power to relieve pain.

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

IV.

DISEASES OF WOMEN.

Sleeplessness in Uterine Disorders.

When patients complain of nervousness or of sleeplessness, the potassic bromide must be given, either alone or in combination with other remedies. A cheap mixture, much thought of by our patients at the University clinic, is the following:

℞ Pulv. ferri sulphat. exsiccat.....gr. xxx.
Potassi bromidi..... } aa ℥ j.
Rad. calumbæ contus..... }
Aquæ bullientis.....Oj.

Steep for twenty-four hours and then strain.

Sig.—One tablespoonful in a wineglassful of water just before or after each meal.

I cannot say much for the palatableness of this infusion nor more for its pharmaceutical elegance; but it does good, and we therefore give it largely to our poor patients. The iron and the potash in it may be increased or lessened, or the former may be left out, as the case may be. The zinc valerianate given thrice daily in doses of from two to four grains is one of our best nervines. For a better class of patients the fol-

lowing antispasmodic mixture can be prescribed with very general satisfaction:

℞ Elixir humuli.....fl. ℥ j.
Elixir ammoniæ valerianat..... } aa fl. ℥ ss.
Syrupi lactucarii..... }

M. Sig.—One dessertspoonful at bedtime, or during the day when needful. *Wm. Goodell, M.D.*

In Rheumatic Dysmenorrhœa.

℞ Am. hydrochlor,..... ℥ iij.
Tinct. stramonii..... ℥ ss.
Tinct. cimicifugæ rac..... ℥ jss.
Syr. glycyrrhizæ..... ʒ ij.

M. Sig.—Teaspoonful three times a day.

Dr. N. S. Davis.

OR

℞ Acidi salicylici..... ʒ iij.
Sodii bicarbonatis..... ʒ ij.
Tinct. stramonii.....
Vini colchici radicis.....aa ʒ iv.
Glycerinæ..... ℥ j.
Aquæ..... ℥ iij.

M. Sig.—A teaspoonful four times a day in water

Dr. N. S. Davis.

When the pain and soreness extends to the ovaries, Professor Davis has succeeded with these formulæ in a large number of cases. He also recommends that patients place themselves in the knee and chest position three or four times a day for a few minutes, thus throwing the uterus, by force of gravity, into its natural position.

Chronic Cervical Endometritis.

℞ Magnesiæ sulphatis..... ℥ ij.
Ferri sulphatis.....gr. xvj.
Acid. sulph. dil..... ʒ j.
Aquæ .....Oj.

M. Sig.—Two tablespoonfuls in a tumbler of iced water daily on rising. *Dr. T. G. Thomas.*

OR

℞ Sodii et potass. tart..... ℥ ij.
Vini ferri amari..... ℥ ii.
Acid. tartarici..... ʒ iij.
Aquæ ..... ℥ xiv.

M. Sig.—Two tablespoonfuls as above.
Dr. T. G. Thomas.

Prof. Thomas advises the use of the above prescriptions as tonic and cathartic, relying mainly in these cases on the observance of good hygiene.

Congestive Dysmenorrhœa.

℞ Ext. ergotæ fl..... ʒ vii.
Tinct. gelsem. co..... ʒ j.
Tinct. aconit. rad.....gtts. xvi.

M. Sig.—A teaspoonful every two, three or four hours. *Bartholow.*

Vomiting of Pregnancy.

℞ Cerii oxalatis.....gr. xxiv.
Ext. hyoscyami.....gr. xxxvi.

M. Et. ft. pil. No. xii, Take one twice a day.

A Remedy for the Sick Stomach of Pregnancy.

Dr. Forwood, at a late meeting of the Lancaster, Pa., Medical Society, read an interesting paper on the "Treatment of the Sick-stomach of Pregnancy." His favorite prescription is as follows:

℞ Rad. columbo .....
Rad. zingiber.....aa ℥ ss.
Fol. sennæ..... ʒ j.
Aquæ bullient.....Oj.
Mix. Infus.
Sig.—A wineglassful before each meal.

Vomiting of Pregnancy.

℞ Cupri sulphat.....gr. ij.
Aquæ distillat..... ℥ ss.
M. Sig.—Six drops at a dose. *Bartholow.*

In Chronic Cervicitis and Leucorrhœa.

℞ Iodoformi..... ʒ j.
Acid. tannici..... ℥ j.
M. Sig.—A sufficient quantity to be packed in a dry state around the cervix. *Bartholow.*

In Anæmia with Amenorrhœa.

℞ Arsenic.....gr. j.
Ferri. sulph. exsiccat..... ʒ ss.
Pulv. pip. nig..... ʒ j.
Pil. al. et myrrh..... ʒ j.
M. Div. in. pil. xl.
Sig.—One twice a day after meals.

Fothergill.

This excellent combination, by its tonic effect on the blood and its determinative effect on the menstrual function, will speedily restore the patient to her normal condition.

The Vomiting of Pregnancy.

℞ Cerii oxalat.....gr. j.
Ipecacuanhæ.....gr. j.
Creasoti..... .....gtt. ij.
Sig.—To be taken every hour. *Dr. Goodell.*

This prescription has been used with uniform success in this annoying and sometimes dangerous complication of pregnancy, at the Episcopal Hospital, Philadelphia.

Milk Fever.

℞ Tinct. aconit. rad.....,.....gtt. xx.
Antimon. et potass. tart.....gr. ij.
Spt. etheris nitrici.....
Syrupi simplicis.....aa ℥ j.
Aquæ aurantii flor..... ℥ ij.
Sig.—A teaspoonful in a wineglassful of sugar and water every two hours. *Dr. Fordyce Barker.*

In addition to this, Dr. Barker gives nutritious, easily digested food; applies the child to the breast as soon as the patient has recovered from the exhaustion of labor; attends to the state of the bowels; gives a diaphoretic anodyne, and has the nurse rub the breasts from circumference to nipple with warm sweet oil every three hours.

In Rheumatic Dysmenorrhœa.

℞ Tinct. cimicifugæ..... ℥ iij.
Tinct. stramonii..... ℥ ss.
Vin. colchici rad..... ℥ ss.
M. Sig.—A teaspoonful at each mealtime in water.

N. S. Davis, M.D.

If the cimicifuga or colchicum causes headache and disturbance of the bowels, lessen the quantity relatively to the other constituents of the prescription. The above is recommended by Prof. Davis in that common class of cases, where pain is severe and the flow scanty.

Laxatives for Habitual Use in Uterine Disorders.

℞ Ext. colcynth. comp.....gr. ij.
Ext. belladonnæ.....gr. ½.
Ext. gentianæ.....gr. j.
Ol. carui.....gtt. ss.
Et ft. pil. No. j.
M. To be taken at bedtime.

The pulvis glycyrrhizæ comp. of the Prussian Pharmacopœia is another good laxative. I have kept patients upon it for months, and always with benefit. The formula for it is as follows:

℞ Pulv. glycyrrh. rad..... } aa ℥ ss.
Pulv. sennæ..... }
Sulphuris sublim..... } aa ʒ ij.
Pulv. fœniculi..... }
Sacchar. purif..... ℥ jss.
M. Sig.—One teaspoonful in half a cupful of water at bedtime.

Wm. Goodell, M.D., Phila.

Metrorrhagia.

℞ Liquor ferri sulphatis..... ℥ ss.
Aquæ..... ℥ j.

M. Sig.—Saturate with this solution the finest cotton wool; dry and expose to moderate pressure. Wrap a sufficient quantity around a long small piece of whalebone and introducing it into the cavity of the uterus, withdraw the whalebone, leaving the cotton *in situ.* Strong thread may be attached to the cotton to withdraw it when necessary. From 12 to 24 hours is as long as it should remain.

This hemostatic is that proposed and used by the distinguished gynæcologist, Dr. J. Marion Sims, which is proof positive of its efficiency.

In Accidental Hemorrhage During Pregnancy.

℞ Acidi sulphurici dil.. ..... ʒ j.
Tinct. opii.....m. xl.
Infus. rosæ. co..... ℥ vj.

Sig.—Two teaspoonfuls every hour.

OR

℞ Plumbi acetatis.....gr. xviij.
Acidi acetici.....m. xx.
Morphiæ acetatis.....gr. j.
Aquæ destillatæ..... ℥ vj.

J. G. Swane, M.D.

Coming from the accoucheur to the Bristol General Hospital, England, these formulæ so uniformly successful may be highly commended. Posture should be attended to and cold compresses applied to the vulva.

Barker's Post Partum Pills.

℞ Ext. colocynth. co.....
Hydr. chloridi mitis.....aa ʒ xiij.
Ext. hyoscyami.....gr. xl.
Ext. nucis vom .....
Pulv. aloes.....
Pulv. ipecac.....aa gr. xx.
Mix. Divide into 120 pills. *Dr. Fordyce Barker.*

For the Anæmia of Chlorosis.

℞ Ferri vini amari..... ℥ vijss.
Tinct. nucis vomicæ..... ʒ iv.
Liq. potassæ arsenit..... ʒ ij.
M. Sig.—A dessertspoonful in a glassful of water just after each meal.

Prof. T. Gaillard Thomas, M.D.

In addition to this Dr. Thomas (regarding the indications to remove the cause, cure the neurosis, and repair the damage) advises general tonic treatment and the observance of good hygiene.

In Menorrhagia of Climacteric Period.

℞ Ext. ergotæ aq. (Squibb)..... ℈ ij.
Butyri cocoæ..... ʒ j.
M. Make twelve suppositories and introduce one into the rectum morning, noon, and night, carrying them well up into the bowel and having the patient lie down for an hour afterward.

Prof. Fordyce Barker, M.D.

The Doctor advises the uses of these suppositories for a week previous to the return of the expected period.

Ergot and Aloes for Amenorrhœa from Atony of the Uterus.

℞ Tr. ergot..... ʒ ij.
Decoct. aloes comp.....ad ℥ viij.
M. Two tablespoonfuls twice a day.

Metrorrhagia of Climacteric Period.

℞ Iodoformi..... ʒ ijss.
Gum tragacanthæ.....gr. xv.
Mucilaginis.....q. s.
M. Sig.—Divide into ten cylinders, each one and one half inches in length, and for the five or six days preceding menstruation introduce one of these cylinders into the cavity of the uterus and keep it in place by a pledget of cotton pushed against the cervix.

Prof. Fordyce Barker, M.D.

This plan Dr. Barker has found to effect a cure if carried out for two menstrual periods.

In Chlorosis.

℞ Ferri arseniat.....gr. ij.
Ext. cinchonæ.....gr. xij.
M. Ft. pil. No. xij.
Sig.—One three times a day after meals.

Dr. Bartholow.

Metrorrhagia.

℞ Hydrarg. chlorid coros.....gr. $\frac{1}{12}$
Tinct. cinchonæ comp..... ʒ j.
Sig.—This amount thrice daily.

Dr. Wm. H. Byford.

The above should be given in the intermenstrual period while mechanical means are resorted to at the time of menstruation. Of course quiet and the best possible hygienic surroundings should be insisted upon.

In Painful Affections of the Breast or Abdomen of Women.

℞ Atropiæ sulph.....gr. iv.
Sp. rectif.....f ℥ ss.
Aquæ destil.....ad. ℥ iv.

Misce ut fiat lotio. (To be applied on lint, covered with oilskin or parydor.) *Dr. Tilt.*

In Nausea of Uterine Irritation.

℞ Liq. bismuthi et ammon. citrat..... ℥ ss.
Acid. hydrocyan. dil.....♏ xxiv.
Sodæ bicarb..... ʒ j.
Tinct. lavand. co..... ..... ʒ iij.
Aquæ, ...f ℥ vj.

Fiat mistura. Cap. cochl. maj. ij. bis die.

Dr. Tyler Smith.

In Obstinate Dysmenorrhœa.

℞ Chloral hydrat..... ʒ j.
Sp. etheris..... ʒ ij.
Liq. opii sedativi..... ʒ ss.
Tinct. hyoscyami..... ʒ iij.
Sp. chloroformi.... ..... ʒ ij.
Aquæ.....ad. ℥ vj.

Fiat mistura. A tablespoonful every two hours.

Dr. Lombe Atthill.

As an Anæsthetic During Labor.

℞ Chloral hydrat..... ʒ iss.
Aquæ aurantii floris..... ℥ vj.

Misce. A sixth part at completion of first stage of labor; another dose in about twenty minutes; and again in about an hour. *Dr. W. S. Playfair.*

In Dysmenorrhœa.

℞ Succi conii..... ʒ j.
Potass. bromid.....gr. xv.
Aquæ.....ad. ℥ j.

Fiat haustus. To be taken three times a day.

Dr. Alfred Meadows.

In Ulceration of Cervix Uteri.

℞ Acidi benzoici.....gr. x.
Acidi tannici.....gr. xv.
Collodion..... ʒ iv.
Acidi carbolici.....gr. xxv.

Fiat lotio. To be applied to affected parts.

Dr. Lombe Atthill.

In Pruritus Vulvæ.

℞ Acidi carbolici..... .gr. x.
Morphiæ acet.....gr. viij.
Acidi hydrocyan. dil..... ʒ ij.
Glycerini..... ʒ iv.
Aquæ.....ad. ℥ iv.

Fiat lotio. *Dr. Lombe Atthill.*

In Menorrhagia.

℞ Acidi gallici.....
Pulv. ergotæ.....ana gr. x.
Fiat pulvis. To be taken every three hours.

Dr. Lombe Atthill.

As a Laxative on Change of Life.

℞ Sulphur. sublim..... ℥ j.
Sodæ bicarb..... ʒ j.
Pulv. ipecac.....gr. v.
Misce. Sumat ℈ j.—℈ ij. h. s. ex lacte.

Dr. Tilt.

Sedative Application in Uterine Affections.

℞ Aconitiæ.....gr. ij.
Atropiæ sulph.....gr. viij.
Morphiæ sulph.....gr. xvj.
Acid. sulph. dil..... ♏ v.
Sp. vini rect..... ℥ ss.
Glycerini..... ℥ ij.
Aquæ.....ad. ℥ iv.
M. Fiat linimentum. *Dr. Tilt.*

Dysmenorrhœa.

℞ Ext. conii.....℈ j.
Ext. strammon.....gr. v.
Ext. opii.....gr. v.
M. Ft. pil. No. x.
Sig.—One pill twice or even three times a day.

Garrigues.

Vaginitis.

℞ Acetanilid.....gr. lxxv.
Ac. tannici.....gr. viij.
Ext. hyoscyami.....grs. j.
Sacchari lactis.....gr. cl.

M. Sig.—Use such a suppository two or three times a day.

OR

℞ Ex. hydrastis fl.....℥ iv.

Sig.—Apply to the cervix and vagina, and place a tampon smeared with vaseline between the vulvæ and in the vagina. *Mundé.*

Vaginismus.

℞ Thymol.....gr. iij.
Ext. belladonnæ.....gr. vj.
Potassii bromid.....ʒ ss.
Ol. theobromæ.....℥ ss.

M. Ft. suppositoria No. iv.

Sig.—One suppository as needed. *De Sintey.*

Amenorrhœa.

℞ Tr. ferri chlor.....f ʒ iij.
Tr. cantharidis.....f ʒ j.
Tr. guaiac ammon.....f ℥ iss.
Tr. Aloes.....f ℥ ss.
Syrupi.....q. s. ad f ℥ vj.

Sig.—Tablespoonful three times a day. *Dewees.*

Vomiting of Pregnancy.

℞ Liq. potass. arsenitis.....℥ ss.

Sig.—Half drop every half hour for six or eight doses.
A. A. Smith.

Chloasma of Pregnancy.

℞ Zinci oxidi.. .....gr. iij.
Hydrarg. ammoniat.....gr. iss.
Ol. theobromæ..... .....
Ol. ricini.....aa ʒ iiss.
Essent. rosæ.....gtt. x.
M. Sig.—Apply to the face night and morning.

Monin.

Metritis—Uterine Tumor.

℞ Tr. aconit. rad.....gtt. xvj.
Ex. gelsemii fl.....ʒ j.
Ex. ergotæ fl.....ad ℥ j.
M. Sig.—Teaspoonful every two to six hours.

Bartholow.

Neurasthenia.

℞ Sodii bromidi.....℥ j.
Tr. nucis vom.....ʒ ij.
Ess. pepsin.....℥ ij.
Aq. menth. pip.....℥ j.
M. Sig.—Teaspoonful before meals and at bed time.

Dr. A. J. C. Skene.

Irritable Bladder.

℞ Lithiæ carbonatis..... .....ʒ ij.
Ac. benzoici.....ʒ iij.
Sodii boratis.....ʒ j.
Aq. dest.....℥ iv.
M. Sig.—A tablespoonful in a glass of water after meals for irritable bladder in women associated with a lithic-acid diathesis. *Dr. Skene.*

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

V.

DISEASES OF CHILDREN.

Diarrhœal Mixture for Children.

℞ Pulv. cret. aromat. (Br)..... ʒ j.
Tinct. opii..... ʒ j.
Syr. zingiberis..... ℥ j.
Decoc. hæmatoxyli.....ad. ℥ vj.
Take two tablespoonfuls after each liquid stool.

Farquharson.

Acute Bronchitis in Children.

℞ Tinct. veratri viridis.....m. xij.
Syr. scilla compos..... ʒ ij.
Syr. balsami tolutani..... ʒ xiv.
M. Sig.—One teaspoonful every two or three hours to a child five years old in the first stages of the disease. *Prof. J. L. Smith, M.D.*

When the temperature falls and the moisture appears on the skin under the influence of the above prescription, we may cease its administration and resort to expectorant mixtures as the following:

℞ Ext. cubeb fl.....m. xl.—ʒ j.
Syr. simplicis..... ℥ ijss.
Sig.—A teaspoonful three or four times a day.

In Summer Diarrhœa of Children.

℞ Plumbi acetat.....gr. viij.
Acid acet.....gtt. vj.
Tinct. opii deodorat.....gtt. iv.
Aquæ destillat..... ℥ j.

M. Sig.—A teaspoonful every two or three hours to a child of two years. *Dr. Roberts Bartholow.*

Infantile Syphilis.

℞ Hydrarg. bichlorid.....gr. j.
Potass. iodidi..... ʒ iv.
Syr. aurantii.....
Aquæ.....aa ʒ ij.

M. Sig.—Five drops for a child about two months old, increased to fifteen or twenty drops if the disease does not yield. *R. W. Taylor, M.D.*

Dr. Taylor has made this subject his especial study, and his experience has been that the above formula is highly efficacious. It is important to suspend the medicine altogether, from time to time, as the system acquires a tolerance for it.

Nocturnal Incontinence of Children.

℞ Strychniæ.....gr. j.
Pulv. cantharides.....gr. ij.
Morph. sulph.....gr. iss.
Ferri pulv..... ℈ j.

M. Ft. pil. No. xl. Sig.—One three times a day to a child ten years old. *Prof. S. D. Gross, M.D.*

This prescription will speedily relieve the irritability of the bladder, especially if conjoined with such

means as a cold shower bath daily, the avoidance of irritant food and late suppers, the patient lying on the side or belly, and taking care to drink nothing for the few hours preceding sleep, and to empty the bladder on going to bed.

Alkaline Mixture for Nursing Children.

℞ Pot. carbonat.....gr. ij.
Ol. cajeput.....m. j.
Aq. anethi..... ʒ ij.
M. Sig.—Three or four times a day. *Fothergill.*

This alkaline mixture will afford speedy relief to infants who have griping pains after taking milk; a condition very frequent and exceedingly annoying.

Catarrhal Bronchitis of Infants.

℞ Syr. allii..... ℥ j.
Syr. ipecac..... ʒ iij.
Syr. tolutani..... ..... ʒ v.
Syr. acaciæ..... ʒ vi.
Tinct. opii camphorat..... ...ad. ℥ iij.
M. Sig.—M. xx. to ʒ j. as needed.

Dr. Farquharson.

Cough Mixture for Infants.

Infants' Hospital, Randall's Island, N. Y.

℞ Tr. opii camph.....
Spts. ammon. arom.....aa fl. ℥ j.
Ext. ipecac fl.....fl. ʒ ss.
Syr. pruni virg.....fl. ℥ j.
Aquæ q. s. ad.....fl. ℥ iij.
M. Dose: a teaspoonful.

Cholera Infantum.

℞ Argenti nitrat.....gr. j.
Acid nitric. dil.....m. viij.
Tinct. opii deodorat.....m. viij.
Mucil. acaciæ..... ℥ ss.
Syr. simplicis..... ℥ ss.
Aq. cinnamoni..... ℥ j.

M. Sig.—A teaspoonful every three, four, or six hours to a child one year old. *Bartholow.*

This combination is remarkably beneficial after the acute symptoms have subsided.

Infantile Convulsions.

℞ Olei succini rectificati.....
Tinct. opii. .....aa ℥ ss.
Olei olivæ..... . .....
Spiritus vini gallici..... ...aa ℥ ij.

Ft. lotio. Rub along the spine.

Dr. Joseph Parrish.

The above will relieve infantile convulsions, and is also an excellent application to relieve the spasms of whooping-cough. Care should be taken to wash the skin with warm water and soap before rubbing in the lotion, so as to promote absorption.

Cholera Infantum.

℞ Cupri sulph.....gr. j.
Tinct. opii deodorat.....gtt. viij.
Aquæ destillat..... ℥ iv.

M. Sig.—A teaspoonful every two, three, or four hours. *Dr. Bartholow.*

Anthelmintic for Round Worm and Ascaris Vermicularis.

℞ Fl. ext. spigeliæ..... ℥ j.
Fl. ext. sennæ..... ℥ ss.
M. Sig.—A teaspoonful to a child of three to five years; or

℞ Fluid ext. spigeliæ et sennæ..... ℥ j.
Santonini.....gr. viij.
M. Sig.—Teaspoonful to a child of five.

Prof. J. L. Smith, M.D.

For the expulsion of the round worm no better formulæ than these have been devised. They are found an effectual means also of destroying the ascaris vermicularis.

In the Delirium of Febrile Affections in Children.

℞ Pot. bromidi.....gr. x.
Chloral hydrat.....gr. v.
Aquæ anethi..... ʒ ij.
M. Sig.—Every six hours, for child of six.

Fothergill.

This combination of bromide of potassium and chloral acts admirably in the cerebral excitement accompanying the evanescent fevers of children.

Children's Diarrhœa.

℞ Infus. rhei..... ℥ ij.
Potass. bicarb..... ʒ j.
Tinct. cinnamoni..... ʒ ij.
Syrupi simplicis..... ʒ vj.
Dose. Teaspoonful every two hours.

Farquharson

In Scarlatina.

℞ Pot. chlorat.....gr. x.
Tinc. ferri perchlor.....m. v.
Syr. zingiberis.....ʒ j.
Aquæ.....ad ℥ ss.

M. Sig.—Every four or six hours for a child from eight to ten years. *Fothergill.*

Dr. Fothergill says of the above prescription that in the severe sore throat of scarlatina he has found it of all others give the best and most satisfactory results.

Bronchitis in Children.

A large sinapism or turpentine stupe, followed by linseed-meal poultices properly made and frequently changed, or a piece of spongio-piline soaked in warm water, are good external applications. The child should be kept in bed. Internally, in acute cases, a dose of calomel with or without jalap may be given at the outset, and expectoration should be assisted by ipecacuanha, squill, citrate of potash, senega, and similar remedies. A useful prescription for a child two to four years old is:

℞ Pot. citrat.....ʒ j.
Vin. ipecac.....ʒ j—ʒ ij.
Tinct. camph. co.....ʒ j.
Syrup. tolutan.....℥ ss.
Aquæ ad.....℥ iij.

Ft. mist. Dose: one teaspoonful every hour or two.

When the secretion becomes abundant, it should be removed from the loaded bronchi by emetics. Mucous and subcrepitant râles are in young children the

best indications for emetics, according to Bouchut. Powdered ipecacuanha, mustard, alum, or sulphate of zinc are best; Vin. Ipecac. sometimes fails even in large doses. A warm bath towards evening is good practice, and the child often sleeps after it, but chill must be carefully guarded against. If there be much restlessness, small doses of Dover's powder at bed-time are valuable. Should pulmonary collapse take place, stimulants, especially Ammon. Sesquicarb., will be needed, with wine and rubefacient liniments. The diet, at first low, must be improved as the disease progresses, care being taken to eliminate all heavy and indigestible things from what is sanctioned. In chronic bronchitis, where there is less fever, and when the child is already exhausted by illness, the cough will require controlling by such drugs as bromide of potassium, belladonna, morphia, etc. Antispasmodics and anodynes, in fact, take the place of expectorants and stimulants. Sinapisms are still useful, and so are stimulating embrocations. Inhalations, too, of steam or medicated vapors (especially that of creasote) are serviceable. The diet must be light and nourishing, and calculated to restrain rather than promote secretion, and therefore especially limited and defined in the matter of fluids.

In such cases quinine is useful in small doses; or if it disagree, Dr. Meigs recommends the following:

℞ Elix. cinchon. flav..... ʒ ij.
Curacoa..... ʒ ij.
Acidi sulph. dil.....m. xij
Aquæ.... ..... ʒ iiss.
Ft. mist., ʒ j. secund. hor.

In very chronic cases, attended with persistent mucous râles over the bases of the lungs, astringents, especially gallic and tannic acids, are useful. I have certainly seen benefit from their employment. At the same time some external agency should not be neglected, as painting with iodine paint or gentle frictions with some stimulating liniment. Dr. Stierlin, of Schaffhausen, recommends carbonate of ammonia rather than emetic or other treatment, especially in the broncho-pneumonia of young children and in the catarrhal attacks of infants. The dose may be up to 5, 10 or even 20 grains. Dr. Stierlin by this means lost only 7 out of 150 cases, whereas Rilliet and Barthez consider the catarrhal pneumonia of infants especially to be almost invariably fatal.

It is convenient here to mention specially another plan of treatment, referred to in former editions, as much in vogue in America and on the Continent, but which, since then, has daily gained ground in England and elsewhere, and which is applicable not only to bronchitis, but to most, if not all, acute inflammations, viz., the treatment by the great vascular sedatives, aconite and veratrum viride.

This treatment, according to the evidence of those who have most largely employed it, is most suitable for children over three years of age, whose previous health has been good, and in whom the inflammation is acute and primary. It is a remark of Bouchut's that in the first stage of childhood the material lesions are less purely inflammatory than in the second stage, and the suppuration of the tissues in both less frequent and of a less laudable quality. It is accordingly found

that these powerful antiphlogistics are of less service during the earliest years of life. Besides the age of the child, the period of the administration of these drugs is an important point; they should be given *as early as possible* in the course of the disease, in *small* and *frequently repeated* doses, until the activity of the inflammation begins to subside, the pulse lowers, the temperature falls, and moisture appears upon the skin. Beyond this point it is unwise to push the remedy, as the depression so induced may be very considerable. Dr. Lewis Smith, of New York, recommends the following prescription for a child five years old in the first stage of acute bronchitis:

℞ Tinct. verat. virid.....m. xij.
Syrupi scillæ comp..... ʒ ij.
Syrupi bals. tolutan..... ʒ xiv.

Misce. One teaspoonful every two to four hours; the medicine to be omitted, or given at a longer interval, if the frequency of the pulse is reduced.

I have but little experience of green hellebore, but I am more and more satitfied that aconite is a most valuable agent when similarly employed. The dose of the tincture of the Ph. B. may be half a drop to a drop, repeated every hour or half hour until the effect described is manifested. The dose is suitable for a child five years old. When the inflammatory symptoms have abated, the ordinary treatment of the special inflammation and its sequelæ is to be resumed; for example, in bronchitis, expectorant mixtures; in tonsillitis, astringent gargles, and so forth. In fact, aconite and green hellebore thus employed may be

regarded as in a measure replacing the depletions and blisterings of days gone by.

Lastly, I think it always desirable, after children have suffered from bronchitis, broncho-pneumonia, etc., that a course of cod-liver oil or the hypophosphites of lime and soda should be advised for a month or two. *E. Ellis.*

In Chronic Diarrhœa and Passive Hemorrhage.

℞ Alumnis.....gr. iij.—xv.
Acidi sulph. dil.....m. ij.—x.
Syrupi.....q. s.
Inf. rosæ acid..... ℥ j.—iv.

To be given every three or four hours. *E. Ellis.*

In the Second Stage of Hooping-Cough.

℞ Ext. conii.....gr. xij.
Aluminis.....gr. xxv.
Aquæ anethi.....f ℥ iij.
Syr. rhœados.....f ℥ ij.

Fiat mistura: capiat cochleare medium sexta quaque horâ. (For a child two or three years old.)
Dr. G. Bird.

In Hooping-Cough.

℞ Cocci pulv.....gr. xij.
Potass. carb..... ℥ j.
Syr. simplicis..... ℥ j.
Aquæ destil..... ℥ iij.

Misce. A teaspoonful when the attack threatens.
Niemeyer.

In Chronic Diarrhœa of Children.

℞ Argenti nitr.....gr. $\frac{1}{6}$ ad $\frac{1}{3}$
Aquæ destil.....℥ j.
Syr. simpl.....ʒ v.

Fiat mistura. To be taken by spoonfuls in the course of the day. *M. Trousseau.*

In Laryngismus Stridulus.

℞ Assafœtidæ.....ʒ ss.
Vitell. ovi No. j. Tere simul, et adde.....
Infus. valerian.....℥ ss.—℥ iv.

For two enemas. *Niemeyer.*

In Hooping-Cough.

℞ Antim. tart.....gr. j.
Tinct. opii.....♏ xx.
Aquæ destil.....f ℥ j.

Misce. A teaspoonful every, or every alternate, night. *Sir. T. Watson.*

Itch in Children.

Sulphur ointment is the general remedy for this disease. The patient having been thoroughly washed, should be rubbed over with it wherever spots exist. If there be unusual thickness of the cuticle, the ointment may be

℞ Adipis.....℥ j.
Sulphur. præcip.....ʒ ij.
Pot. bicarb.....ʒ j.

Ft. unguent.

A strong alcoholic solution of stavesacre is often efficacious, or the Pulv. staphisagriæ may be combined with Ung. Sulph. The oil of chamomile is also stated to be useful in the Ung. Sulph. A lotion of pentasulphide of calcium is recommended by Mr. Erasmus Wilson. If it be desired to conceal the fact of using sulphur ointment, it may be colored with Hyd. Bisulph (Cinnabar), and scented with oil of bergamot. Styrax is also occasionally employed, *e. g.*:

℞ Styracis liquid..... ℥ j.
Sp. rectif..... ... ʒ ij.
Ol. olivæ..... ʒ j.
Ft. unguent.

Other formulæ are:

℞ Acid carbolici..... ʒ ij.
Glycerini..... ℥ ij.
Ft. applicatio.

℞ Hyd. perchlor.....gr. iv.
Sp. vin. rect..... ʒ vj.
Ammon. hydrochlor..... ʒ ss.
Aq. rosæ.....ad ℥ vj.
Ft. lotio.

℞ Potassii sulphureti..... ℥ vj.
Sapon. alb.....lb. ij.
Ol. olivæ.....Oij.
Ol. thymi..... ʒ ij.
Ft. applicatio.

This is a powerful preparation, and sufficient for inunction from top to toe (as is often required) for several days. *E. Ellis.*

In Diarrhœa.

℞ Cretæ preparatæ..... ʒ iss.
Acaciæ pulv.....
Sacchari albi.....aa ʒ j.
Tinct. opii.....m. x.
Aquæ..... ℥ iij.
Ft. mist. A teaspoonful every hour. *Dewees.*

In the Irritable Stomach of Young Children, Accompanied with Vomiting, of Sour and Curdled Character.

℞ Pot. bromid.....gr. j.—iij.
Mist. cretæ..... ʒ j.—ij.
Syrupi.....q. s.
E. Ellis.

In Hooping-Cough.

℞ Aluminis.....gr. xxiv.
Acidi sulph. dil.....♏ xij.
Syr. rhœados.....f ʒ iv.
Aquæ.....f ℥ iis.
Fiat mistura: sumat f ʒ iij. sextis horis.
Dr. West.

In Simple Diarrhœa of Infants.

℞ Magnes. sulph..... ʒ j.
Tinct. rhei..... ʒ ij.
Aquæ carui..... ʒ vij.
Misce. A teaspoonful every six hours.
Dr. West.

As a Stimulant in Croup.

℞ Etheris acet..... ʒ iij.
Camphoræ.....gr. x.

Misce. Ten to fifteen drops every quarter of an hour. *Niemeyer.*

In Porrigo of Children.

℞ Acidi acetici diluti.....
Mel. boracis.....āā ℥ j.
Aquæ..... ℥ iv.

Misce. Fiat lotio. *Dr. Headland.*

In Severe Coughs of Children of Four Years of Age.

℞ Tinct. camphoræ co.....♏ xx.
Vini ipecacuanhæ.....♏ x.
Vini antim.....♏ xxx.
Mucil. acaciæ..... ʒ vij.

Misce. Sumat ℥ ij. quartâ quâque horâ.
Dr. West.

In Hooping-Cough.

℞ Tinct. opii..... ʒ ij.
Tinct. cantharidis..... ʒ ij.
Lin. camphoræ co..... ℥ j.

Misce. Fiat linimentum. To be rubbed on the chest. *Dr. West.*

In Chorea.

℞ Liq. arsenicalis.....♏ x.

To be taken immediately after meals. For child 5 to 12 years. *Dr. Eustace Smith.*

In Febrile Coughs of Children One Year Old.

℞ Potass. bicarb.....
Acidi citrici.....ana ℈ j.
Syr. mori..... ʒ iij.
Aquæ destil..... ʒ ix.
Misce. Sumat ʒ ij. sextâ quaque horâ.

Dr. West.

An Aperient for Children of Three Years of Age.

℞ Potass. sulph.....gr. xij.
Inf. rhei..... ʒ vss.
Tinct. aurantii..... ʒ ss.
Aquæ cinnam..... ʒ iss.
Misce. Capiat ℥ ss pro re natâ. *Dr. West.*

In Scarlatina.

℞ Solut. chlori. (Midd. Hosp.)..... ʒ ij.
Aquæ.....Oj.
Misce. Sumat cochl. j. ad ij. frequenter in die pro-ratione ætatis. *Sir T. Watson.*

Stomatitis.

These cases are usually easily cured by the use of chlorate of potash in good doses, given three or four times a day, the mouth being rinsed well with a weak solution of the chlorate, or in young children syringed therewith, if the child be too young to rinse the mouth ; glycerine of borax may be applied after each cleansing. The stomach and bowels must be regulated; and when improvement takes place, tincture or decoction of bark will be of great use—in fact, the

best form of tonic. Nutritious diet and wine are often necessary, when the child is low.

Chloride of lime is used by M. Bouchut, forty-five grains of the chloride to six drachms of honey, and application made with a camel-hair brush. Dilute nitric and hydrochloric acids are occasionally valuable in obstinate cases as local applications ; acid and bark or ammonia and bark being given meanwhile internally. Dr. Dewees especially recommends:

℞ Cupri sulph.....gr. x.
Pulv. cinchon. opt.....ʒ ij.
Pulv. gum. Arab.....ʒ j.
Mel. commun.....ʒ ij.
Aq. fontan.....℥ iij.

Ft. applicatio. The ulcerations to be touched twice a day with it. *E. Ellis.*

Cretinism.

℞ Ext. thyroid.....℥ j.

Sig.—Begin with one-half grain twice daily, increasing to one grain or one and a half grains or more twice or three times daily. The results of the thyroid treatment of cretinism are among the most brilliant in modern medicine.

Adenitis of Children.

℞ Ichthyol.....
Ung. hydrarg.....
Ung. belladonnæ.....aa ʒ j.
Ung petrolati.....℥ ss.
Ft. ung.....

M. Sig—Apply night and morning over affected glands, using friction till absorbed.

Whooping-Cough.

℞ Antipyrin.....gr. xvj.
Syr. tolutani..... .....
Aquae.....aa ℥ ss.

M. Sig.—A teaspoonful every four hours to a child four years old; the interval may be gradually reduced to two hours. *Dr. F. M. Crandall.*

Holt, Crandall Kerley and others speak of antipyrin as the most effective drug in whooping cough. Holt advises the following doses: at six months the initial dose should be one grain every three hours, the interval being reduced if necessary to two hours; at two years the initial dose should be two grains every four to six hours, the interval being gradually reduced to two hours. Kerley combines bromide of sodium with the antipyrin. Three grains of bromide may be given every three hours at three years.

Bromoform is effective in some cases, but requires care in its use. The following is an available formula:

℞ Bromoformi.....♏xl.
Mucilag. gummi Arabic.....
Syr. tolutani.....aa ʒ vij—M.

Sig.—Teaspoonful from three to five times a day.

Tonic for Children.

℞ Ferri oxid, saccharated.....gr. iij.
"Eisenzucker."

Ft. tablet.

Sig.—One or two after meals for anaemia. A palatable and effective form of iron.

Eczema of Children.

℞ Pulv. calaminæ prepar. ..... ʒ ij.
Zinci oxid. ..... ℥ ss.
Glycerinæ. ..... ℥ ss.
Liq. calcis. ..... ℥ ij.
Aq. rosæ ..... ad ℥ viij.

M. Sig.—During stage of acute inflammation, dip a piece of muslin in this solution and apply with a bandage.

Dr. L. E. Holt.

℞ Ac. salicylici ..... gr. v.
Zinci oxidi ..... ʒ ij.
Amyli ..... ʒ ij.
Vaseline ..... ℥ j.

M. Sig.—" Lassar's paste " for use after the more acute stage has passed. A most useful prescription, especially in eczema behind the ears and in folds of the skin.

Lassar.

℞ Ung. picis liquidæ ..... ʒ iij.
Zinci oxidi ..... ʒ iss.
Ung. aquæ rosæ ..... ʒ vj.

M. Sig.—For use in chronic cases. Keep in close contact with diseased area by spreading upon muslin and applying under a bandage. *Dr. L. E. Holt.*

℞ Pulv. tragacanthæ ..... ʒ j.
Glycerinæ ..... ℥ iss.
Aq rosæ ..... ℥ iv.

M. Sig —" Pick's paste " for use where a mask or ointment cannot be used. To this may be added zinc oxide, carbolic acid, ♏v, or tar, ♏x.

Regulation of the diet is of more importance in eczema of children than in that of adults. Each case, however, must be studied and treated individually. In infants, reduction of the fat is often followed by improvement. In older children, on the other hand, who are thin and poorly nourished, the administration of cod-liver oil and iron is often equally efficacious. Rhubarb and soda is almost invariably useful.

Chorea.

℞ Liq. potassii arsenitis.....ʒ j.
Vini ferri amari..... .....ʒ iij.

M. Sig.—Begin with fifteen drops, three times a day, and increase one drop at each dose until the point of tolerance is reached. Reduce the dose to one half and again slowly increase to a point a little below that first reached and then continue the administration until improvement is seen. *New York Polyclinic.*

Enuresis.

℞ Atropiae sulphatis.....gr. ss.
Aquae destillatae.....℥ j.

M. Sig.—One drop for each year of age, given at four and seven o'clock in the evening.

The value of atropin in full doses cannot be doubted. The urine of the enuresis patient should always be examined and the treatment instituted which the results of such an examination indicate. A permanent cure cannot always be effected until constitutional faults are remedied.

Pleurisy.

℞ Spt. ætheris nitrosi.....gtt. xx.
Liq. ammon. acet.....ʒ ss.
Chloroformi.....gtt. ij.
Aq. menthæ vir.....q. s. ad ℥ j.

M. Sig.—One dose every two hours for a child of four years.

For Thrush.

℞ Sodii bicarbonatis.....gr. v.
Sodii boratis.....gr. v.
Aquæ.....℥ j.

M. Sig.—Mouth wash to be applied four times a day upon a soft cloth wrapped upon the finger or by a cotton swab. *Dr. F. M. Crandall.*

Rickets.

℞ Ol. Morrhuæ.....℥ iv.

Sig.—One to two teaspoonfuls three times a day after meals. If the pure oil is not taken well use an emulsion. (See Section X.)

Although cod-liver oil is the most useful medicinal agent for rickets, it is of less importance than diet. It is well established that the disease is due to deficient fat in the diet and excess of carbo-hydrate, together with absence of sunlight and unhygienic surroundings. Hence, increase the cream in the food as much as the child will digest and withdraw all starches under one year and reduce them over that age. Keep the child in the open air as much as possible. Proper diet alone will cure rickets, but cod-liver oil and tonics will aid.

Tapeworm.

℞ Ol. filicis maris.....ʒ iij.
Ol. chenopodii.....ʒ j.
Ol. terebinth.....ʒ ij.
Emul. ol. ricini (50 per cent.).....q. s. ad ℥ ij.

M. Sig.—Teaspoonful twice a day for a child of six years. *Dr. L. Starr.*

For Fever in Children.

℞ Spts. ætheris nitrosi.....ʒ ij.
Liq. ammonii acetatis.....℥ ss.
Glycerini.....℥ ss.
Aquae cinnamomi.....ad ℥ iij.

M. Sig.—A teaspoonful every two hours to a child of five years. This is an effective and palatable mixture.

Dr. F. M. Crandall.

Laxative Alkaline Mixture.

℞ Sodii bicarbonatis.....ʒ ss.
Syr. rhei aromat.....℥ ss.
Syr. sennæ.....℥ ss.
Syr. amant. cort.....℥ i.

M. Sig.—A teaspoonful, more or less, three times a day at one year. *Dr. J. P. C. Griffith.*

Convulsions.

℞ Chloralis hydratis.....gr. v.
Aquæ.....ʒ j.

M. Sig.—Give by high rectal injection to a child of one year and repeat in one hour if necessary. If the choral is expelled or the convulsions continue give morphine hypodermically, $\frac{1}{24}$ of a grain at one year.

Holt.

Earache.

℞ Atropinæ sulphates.....gr. j.
Glycerini.....
Aq. destil.....aa ʒ j.

M. Sig.—Three or four drops to be warmed and dropped into the ear every three hours. For the earache of scarlet fever or from reflex causes.

Dr. T. M. Rotch.

Summer Diarrhoea.

℞ Bismuthi subnitratis.....gr. xv-xx.
Syr. rhei. aromat.....♏iii.
Aquae.....ad ʒ j.

M. Sig.—This amount to be given every two hours after the bowels have been acted upon by castor oil or calomel. Milk should be stopped absolutely for at least one day, and longer if the bowels continue to be loose and the stools of green color. Castor oil, calomel, bismuth, and opium are the four most useful drugs. Intestinal antiseptics give but little aid, and are dangerous; the new astringents have a very limited field of usefulness. For irrigation of the bowel boric acid solution or normal salt solution (heaping teaspoonful to a pint) is best, the temperature of which should range from 95° to 100°.

Dr. C. G. Kerley.

Asthma in Children.

℞ Antipyrin.....gr. iv.

Sig.—For one dose at bedtime to a child of five years when the attacks are recurrent at night.

Dr. L. E. Holt.

Summer Diarrhœa.

℞ Bismuthi subgallat.....gr. xxiv.
Syr. rhei aromatici..... .ʒ j.
Mist. cretæ.....℥ ij.
Aquae.....ad ℥ iii.

M. Sig.—A teaspoonful every two hours at one year for summer diarrhoea, particularly with flatulence and irritable stomach.

The bowels should first be emptied by means of rhubarb, calomel, or the following:

℞ Syr. rhei..... .....
Ol. ricini.....aa ℥ ss.

M. Sig.—One to three teaspoonfuls according to the age of the child. All milk should be withheld. Opium is indicated as follows: (1) in the first stages of acute diarrhoea, before the intestinal canal has been freed from decomposing matter; (2) when the passages are infrequent or of bad odor; (3) when there is a high temperature or cerebral symptoms are present; (4) when its use is followed by elevation of temperature or the passages become more offensive. It is indicated (1) when the passages are very frequent, with pain; (2) when the passages are excessively frequent, large, and watery; (3) in dysenteric diarrhoea preceded by castor oil or a saline; (4) in late stages, with small, frequent, nagging passages; (5) when the passages consist largely of undigested food, and the bowels act as soon as food is taken into the stomach. The opiate should not be added to the ordinary diarrhoea mixture, which is usually repeated at short intervals. It should be given alone. The dose should be as small as possible, the object being to check

peristalsis and relieve excessive pain. Narcotism should be strictly avoided. The dose of paregoric may be as follows: At three months, μ ii; one year, μ viii; five years, μ xxx. The dose of the deodorized tincture may be at three months, μ $\frac{1}{12}$; one year, μ $\frac{1}{3}$; five years, ♏ ii. The dose of Dover's powder may be at three months, gr. $\frac{1}{12}$; at one year, gr. $\frac{1}{3}$; at five years, gr. ii.

F. M. Crandall.

The Treatment of Diphtheria.

The treatment of diphtheria has been revolutionized since 1895. Of the many drugs used prior to that date, but few are now indicated. The tincture of iron in full doses is valuable for both its local and constitutional action. Stimulants should be used freely to prevent exhaustion and heart failure. Of the various stimulants alcohol is the most effective. Absolute quiet in the recumbent position should be insisted upon. Bland but very mild antiseptic washes are applied to keep the diseased parts cleansed and to wash away the bacilli as they are liberated upon the surface. The important features of the modern treatment of diphtheria are rest, nourishment, stimulants, bland antiseptic washes, and antitoxin.

The local applications in most common use in the hospitals of New York, Boston, and Philadelphia are normal salt solution, saturate solution of boracic acid, Seiler's solution, or a bichloride solution of 1 to 10,000. All of these solutions are used warm. The following is a good bichloride solution rendered milder by the addition of borax and glycerine:

℞ Hydrargyri chloridi corrosivi.....gr. ¾
Sodii boratis ....ʒ ij.
Glycerini.....℥ ij
Aquae.....q. s. ad O j.

M. Sig.—For gargle or throat wash.

The value of the antitoxin treatment of diphtheria has been more certainly established than almost any other method of treatment in the whole field of medicine. Few drugs have been so strongly criticised and still fewer have borne the ordeal so triumphantly. It should be used in all cases of diphtheria, whether mild or severe. It should be used in every suspicious case without waiting for the result of bacterial examination, so important is the value of early administration. This is shown by the report of the American Pediatric Society upon 4,120 cases. The mortality among those receiving the injection on the first day was 4.07; the second day, 7.04; third day, 8.08; fourth day, 20.07; fifth day, 35.03.

The doses are now larger than those at first employed, and the results are correspondingly better. For a child of over two years, in laryngeal and severe cases, 3,000 units should be given, to be repeated in from twelve to twenty-four hours if there is no improvement, a third dose after a similar interval if required. Under two years and for very mild cases 1,500 units should be given. The dosage should always be estimated in units and not in the volume of serum. Only serum produced by well-known firms should be employed.

Nasal diphtheria requires a large dose of antitoxin (not less than 3,000 units) for the condition is always dangerous. The nostrils should be frequently syringed with one

of the solutions already mentioned. Hemorrhage should be stopped by the use of a solution of antipyrin (1 to 5) or suprarenal extract (five grains to one drachm of water).

Croup is now known to be, in the great majority of cases, diphtheritic in nature. The antitoxin treatment has been more effective in croup than in any other form of diphtheria. It prevents the necessity of the intubation tube in a large number of cases and has reduced the mortality in both operative and non-operative cases to a point that would have seemed incredible ten years ago. The dose should be large and should be given promptly at the first signs of laryngeal obstruction. Emetics and the various depressing treatments formerly in use should never be prescribed. Intubation should be performed if the disease is not controlled by antitoxin. Tracheotomy has been entirely supplanted by intubation in croup in America.

The value of "immunizing doses" of antitoxin has been abundantly established. For this purpose from 300 to 500 units are employed in children of three to eight years, and 50 units in infants of three months. Immunity lasts for at least twenty days. Every child in a family where diphtheria develops should be immunized, and every adult who must come in contact with the patient.

F. M. Crandall.

VI.

FEVERS, RHEUMATISM, GOUT, AND MALARIA.

Simple Continued Fever.

℞ Acid hydrobrom..... ʒ j.
Syr. simplicis..... ʒ ij.
Aq .....ad. ℥ j.
M. Sig.—Every hour. *Fothergill.*

Dr. Fothergill, in speaking of the above formula, says it will probably constitute *par excellence* the fever mixture of the future. It is especially indicated where there is cerebral disturbance.

Delirium of Typhoid.

℞ Antim. et potass. tart.....gr. j.—ij.
Morphiæ sulph.....gr. iss.
Aq. lauro-cerasi..... ℥ j.
M. Sig.—A teaspoonful every two, three, or four hours. *Dr. Bartholow.*

This will quiet delirium and produce sleep in the abnormal wakefulness of typhoid. It may also be advantageously used for these same conditions, viz., delirium and wakefulness of delirium tremens.

Local Application in Rheumatic Pains.

℞ Lin. aconit.....ʒ ij.
Lin. belladonnæ.....ʒ ij.
Glycerine.....ad. ℥ ij.

Stg.—Apply locally over seat of pain. *Fothergill.*

This liniment is especially advised when the pain is confined to the sciatic nerve, offering marked relief. Treacle may be substituted for the glycerine with advantage. A convenient mode of application is to spread the above on a V-shaped piece of lint and place on the thigh.

Fevers.

℞ Acid. hydrochlor. dil.....m. xv.
Syr. aurantii.....ʒ j.
Aquæ.....℥ ij.

M. Sig.—Give at a draught to a fever patient every three or four hours. *Dr. Fothergill.*

The above acid drink, combined with fresh air and good general hygiene, will be found of more benefit in most fevers than the more powerful remedies of therapeutics, as was sufficiently demonstrated in the treatment of fevers in the late civil war.

Anti-Rheumatic Pills.

℞ Ext. colocynth co.....gr. 45.
Ext. colchici acet.....gr. 30.
Ext. hyoscyami.....gr. 10.
Hydrarg. chlor. mit.....gr. 10.

Mix. Divide into thirty pills.

In Chronic Malarial Indispositions.

℞ Pil. ferri carbonat..... ʒ j.
Acidi arseniosi.....gr. j.
M. Ft. pil. No. xx. Sig.—One three times a day.

OR

℞ Quiniæ sulph.....℈ ij.
Ferri sulph. exsiccat.....℈ j.
Acidi arseniosi..... .....gr. j.
M. Ft. pil. No. xx. Sig.—One pill three times a day. *Bartholow.*

In Symptomatic Fevers.

℞ Vin. antim.....m. xx.
Tinct. hyoscyam..... ʒ ss.
Liq. ammon. acetat..... ℥ j.
M. Sig.—Every four or six hours. *Fothergill.*

Where there is general disturbance of the functions, the tongue furred, appetite lost, general malaise and weakness, and where the face is flushed, head hot, and painful, the above draught, in connection with a light opiate at night, will yield marked relief.

Sthenic Fever.

℞ Vini antim.....m. xx.
Tinct. opii.....m. xv.
Liq. amm. acetat..... ℥ j.
M. Sig.—Every four or six hours. *Fothergill.*

This combination of opium, antimony, and ammonia will be found very efficient in those fevers characterized by bounding pulse and violent delirium.

In Lithiasis.

℞ Pot. bromidi.....gr. xx.
Tinct. hyoscyami.....ʒ ss.
Tinct. lupuli.....ʒ j.
Mist. camph.....℥ j.
M. Sig.—As a draught at bedtime. *Fothergill.*

In the restlessness and suffering of lithiasis, where sleep must be had, and where opium in any form is to be strictly avoided, the above formula is a very excellent hypnotic. If desired, tincture of cannabis indica (15 drops) may be substituted for hyoscyamus.

In Gout.

℞ Mag. sulph.....ʒ ij.
Pot. bicarb.....gr. xv.
Tinct. sem. colchici.....m. x.
Inf. buchu.....℥ j.
M. Sig.—Every four or six hours. *Fothergill.*

Mercury as a purge and opium as a sedative are often indicated in chronic renal changes, but are ill borne by patients in this condition. As a substitute the above is an excellent remedy. It should be followed by free draughts of water.

In Chronic Rheumatism.

℞ Acid arseniosi.....gr. iij.
Pulv. guaiaci.....ʒ iij.
Pulv. capsici.....ʒ ss.
Pil. al. et myrrh.....ʒ iij.
M. Div. in pil. cxx. Sig.—One pill twice a day.
Fothergill.

Coming from one whose practice was for a long time in a district where chronic rheumatism was very prevalent, this formula is invaluable. It should be combined with general tonic treatment, especially fatty foods.

Diarrhœa of Typhoid.

℞ Argenti nitrat.....gr. iij.
Pulv. opii.....
Pulv. ipecac.....aa gr. vj.

M. Ft. pil. No. xij. Sig.—One every four or six hours. *Bartholow.*

When the diarrhœa of typhoid fever resists bismuth, Hope's mixture and laudanum enemata, the above prescription will often magically relieve. It is also a most effective remedy in the diarrhœa of phthisis.

Malarial Enlargement of the Spleen.

℞ Quiniæ sulphat..... ʒ j.
Ferri sulphat. exsiccat..... ʒ iss.

M. Ft. pil. No. xxx. Sig.—Four or five pills during the day. *Dr. Bartholow.*

Or the following is equally efficient, and is sometimes better borne by the stomach:

℞ Pil. ferri carbonat..... ʒ j.
Acidi arseniosi.....gr. j.
Quiniæ sulph..... ℈ ij.

M. Ft. pil. No. xl. Sig.—Two pills three times a day. *Dr. Bartholow.*

In Acute Fever.

℞ Tinct. opii.....m. xv.
Vin. antimon.....m. xx.
Mist. camph..... ℥ j.
M. Sig.—Every four or six hours. *Fothergill.*

In fevers dependent on inflammatory affections other than those of the brain, where there is often delirium, depending on vascular fulness, this formula will be found of great benefit.

Anæmia of Chronic Malarial Poisoning.

℞ Chinoidin .....ʒ ij.
Resinæ podophylli.....gr. iv.
Ferri sulphat. exsiccat.....ʒ j.
M. Ft. pil. No. xx. Sig.—One three times a day.
Dr. Bartholow.

The use of the above formula is highly commended by Dr. Bartholow, especially where there is enlargement of the spleen and engorgement of the portal circulation.

Acute Rheumatism.

℞ Tinct. ferri..... ℥ ij.
Sig.—Twenty to thirty minims every four hours.
Bartholow.

The above form of iron is more especially adapted to the treatment of pale and delicate cachectic subjects. In these cases it diminishes the pain, fever and sweats, lessens the chances of cardiac mischief, and hastens canvalescence, by retarding waste and favoring excretion of uric acid through the kidneys.

In Rheumatism.

℞ Am. chlor.....gr. xv.
Mist. guaiaci..... ℥ j.
M. Sig.—Three or four times a day. *Fothergill.*

In the rheumatism of young persons, often dependent on or at least associated with inactivity of the skin, this is an excellent formula. When there is a history of exposure, as in old persons, the skin should be covered with flannel, and the following may be given three or four times a day:

℞ Pot. bicarbonat.....gr. x.
Pot. iodidi.....gr. v.
Mist. guaiaci..... ℥ j.

In Hectic Fever.

℞ Tinct. digitalis..... ʒ iij.
Tinct. ferri chloridi..... ʒ v.
M. Sig.—Fifteen drops three or four times a day.
Dr. Bartholow.

This will be found to abate the temperature and diminish the sweats in hectic fever.

In Muscular Rheumatism.

℞ Ammon. muriat..... ℥ j.
Ext. cimicifugæ..... ℥ ij.
Syr. simplicis.....
Aquæ laur. cerasi.....aa ℥ j.
M. Sig.—A teaspoonful three or four times a day.
Bartholow.

Gout.

℞ Tinct. colchici seminis..... ....♏ xx.
Potass. bicarbonat.....gr. x.
Aquæ pimentæ..... ℥ j.
M. Sig.—A draught.

This is the standard formula used at the London Hospital for gout.

That used at the University Hospital is the following:

℞ Tinct. colch. sem.....♏ xv.
Magnes. carb.....gr. vj.
Magnes. sulph.....gr. xxx.
Aq. menth. pip. ad..... ℥ j.

The Rheumatism of Phthisis.

℞ Potass. iodidi..... ʒ j.
Fl. ext. conii..... ʒ iij.
Tr. opii. camph..... ℥ ij.
Aq. aurant. flo..... ʒ iv.
Aquæ..... ℥ iv.
M. Sig.—A teaspoonful thrice daily.

This combination of drugs has been found to exert a markedly favorable influence over the course of this disease. It is the standard formula of the Roosevelt Hospital for these cases.

Fevers.

℞ Quin. sulph.....gr. v.
Tinct. digital..... .....♏ xv.
Ac. phos. dil.....♏ xv.
Aquæ..... ℥ j.
M. Sig.—Every four or six hours. *Fothergill.*

In treating pyretic conditions where the heart is acting feebly and irregularly, and the skin is moist rather than dry, this union of antipyretics with a mineral acid will be very grateful to the patient and highly efficient.

Rheumatism.

℞ Lithii bromidi..... ʒ iij.
Syr. zingiberis..... ℥ ss.
Aquæ..... ℥ iss.
M. Sig.—A teaspooful three times a day.

Bartholow.

Very serviceable where the smaller joints remain swollen after the acute symptoms have subsided.

Salicylic Acid Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Acidi salicylici.....gr. 160
Potass. acetat.....gr. 320
Glycerinæ.....fl. ℥ 1
Aquæ q. s. ad.....fl. ℥ 4
Mix. Dose: a teaspoonful.

Anti-Rheumatic Mixture.

Bellevue Hospital, New York.

℞ Sodii et potass. tart..... ℥ ½
Potass. nitrat..... ʒ 5
Vin. colchici sem.....fl. ʒ 2
Aquæ q. s. ad.....fl. ℥ 2
Mix. Dose: a teaspoonful.

Anti-Rheumatic Mixture.

Hart's Island Hospital, New York.

℞ Potass. iodid ..... ℥ i
" acetat ..... ℥ iv
Tinct. colchici sem ..... fl. ℥ ii
Aquæ ..... O ii
Mix. Dose: a tablespoonful.

Anti-Rheumatic Mixture.

Charity Hospital, Blackwell's Island, N. Y.

℞ Sodii et potass. tart ..... ℥ iss
Vini colchici sem ..... fl. ʒ ii
Aquæ q. s. ad ..... fl. ℥ ii
Mix. Dose: a teaspoonful.

Pulvis Quiniæ Co.

(*Clark's Powder.*)

℞ Quiniæ sulphat ..... gr. x
Pulv. capsici ..... gr. iii
Pulv. opii ..... gr. i
M. Dose for ague. *Dr. Alonzo Clark.*

In Insomnia and Delirium of Typhus Fever.

℞ Liq. opii sed ..... ʒ j.
Antimon. et pot. tart ..... gr. j.
Mist (aquæ) camph ..... gr. vj.
Sig.—Tablespoonful every hour till sleep is induced.

The above is highly commended by Graves and Murchison, who claim for it magical effects, the patient waking refreshed and rational.

White's Gout Pills.

℞ Hydrarg. chlor. mit.....
Pulv. aloes.....
Pulv. ipecac.....
Extr. colchici acet.....
} aa ʒ i

Mix. Divide into 60 pills.

In Typhoid Fever.

℞ Acid. hydrochlor..... ʒ ss.
Syr. simplicis..... ℥ j.
Mucil. salepæ..... ℥ vj.

Misce. A tablespoonful every two hours.

Niemeyer.

In Rheumatic Gout in Cachectic Subjects.

℞ Liq. potass..... ♏ xv.
Potass. iodidi.....gr. iij.
Ext. sarsæ..... ʒ j.
Decoc. cinchonæ..... ʒ xj.

Misce. Fiat haustus, ter in die sumendus.

Dr. H. W. Fuller.

In Chronic Rheumatism of Cachectic Subjects.

℞ Ol. morrhuæ..... ʒ iij.
Liq. potass..... ♏ xx.
Potass. iodidi.....gr. iij.
Ess. limonum..... ♏ viij.
Aquæ destil..... ℥ j.

Misce. Fiat haustus, ter in die sum.

Dr. H. W. Fuller.

In Neuralgic Rheumatism.

℞ Ol. santalini.....♏ ij.
Veratriæ .....gr. xxxvj.
Ol. olivæ.....ʒ ss.
Adipis.....ʒ viiss.
Ol. bergamii.....♏ x.
Misce. Fiat unguentum. *Dr. H. W. Fuller.*

In Asthenic Inflammatory Fever.

℞ Tinct. camph. co.....♏ xx.
Ammon. carb.....gr. v.
Decocti senegæ.....℥ iss.
Fiat haustus. *Mr. Erichsen.*

In Gout.

℞ Ext. colchici acet.....gr. vj.
Ext. rhei.....gr. vj.
Ext. aloes socot.....gr. vj.
Ext. belladonnæ.....gr. j.
Misce. Fiant pil. vj. sumat j nocte, bis hebdomad.
Dr. A. B. Garrod.

In Injuries of Muscles and Tendons, with Rheumatic Diathesis.

℞ Ext. colchici acetici.....gr. j.
Pulv. ipecacuanhæ co.....gr. x.
Ext. colocynth. co.....gr. iv.
Fiant pil. iij. *Mr. Erichsen.*

An Aperient in Gout.

℞ Tinct. colchici sem.....♏ xv.
Mag. carb.....gr. vj.
Sulph. mag.....ʒ ss.
Aquæ menth. pip.....ad ℥ j.
M. Fiat haustus. *University Hospital.*

In Gout.

℞ Pulv. colchici sem.....ʒ ss.
Calomelanos.....gr. viij.
Pulv. digitalis.....
Quin. sulph.....aa gr. xv.
Ext. colocynth.....gr. viij.
Conf. rosæ.....q. s.
M. Ut fiat massa in pil. xx. dividenda. (1—4 during the day, at the commencement of the attack.)
Trousseau and Reveil.

Rheumatism.

℞ Sodii salicylatis.....℥ ss.
Tr. gentianae comp.....℥ j.
Elixir simplicis.....℥ ij.
M. Sig.—A teaspoonful every two to three hours until the symptoms are relieved, when the doses are continued every three or four hours. *Dr. F. M. Crandall.*

No other remedy relieves the pain and fever of rheumatism as quickly and certainly as does salicylic acid and its compounds. The following is another method of administration:

℞ Sodii salicylat.....ʒiss–ʒij.
Syr. aurantii amar. cort.....
Aq. destillat.....aa ℥ ij.
Curacoa.....℥ j.
M. Sig.—To be taken in carbonated water in the course of twenty-four hours.

Liniment for Rheumatism.

℞ Ol. Gaultheriæ .....
Ol. olivæ.....
Lin. saponis.....
Tr. aconiti.....
Tr. opii.....aa ʒij.
Ft. liniment.
M. Sig.—Apply freely and cover with cotton batting.

Chronic Rheumatism.

℞ Potass. iodid.....ʒiij.
Vini colchici sem. .....
Tr. opii camph.....aa ℥ij.
Tr. stramonii.....ʒvj.
Tr. cimicifugæ.....℥iij.
M. Sig.—Teaspoonful three times a day.

St. Luke's Hospital, N. Y.

Gout.

℞ Ex. colchici acetici.....
Ex. aloes.....
Pulv. ipecac.....
Hydrarg. chlor. mitis..... aa gr. j.
Ex. nucis vomicæ.....gr. ¼
M. Sig.—One such pill to be taken every four hours until purgation ensues. *Dr. A. L. Loomis.*

This is a most reliable prescription for relieving the severe pain of a gouty attack.

For the Diarrhea of Typhoid Fever.

℞ Ac. sulphuric. aromat.....ʒj.
Ext. hæmatox. fl..... .....℥ss.
Spts. chlorof..... .....℥ss.
Syr. zingiberis.....ad ℥iij.
M. Sig.—Two teaspoonfuls every two to four hours.

Hare.

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

VII.

VENEREAL DISEASES.

Syrup of Biniodide of Mercury.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Potassii iodidi.....gr. 80
Hydrarg. biniodidi.....gr. 1½
Syrupi.....fl. ℥ 2
Dissolve and mix. Dose: a teaspoonful.

Dr. Banks.

Thompson's Mixed Treatment.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Hydrarg. biniodidi.....gr. 1
Potassii iodidi.....ʒ 3
Tinct. aurantii.....fl. ℥ 1
Aquæ.....fl. ℥ 3
Mix. Dose: a teaspoonful.

Dr. Beverhout Thompson.

Taylor's Mixed Treatment.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Hydrarg. biniodidi.....gr. 1
Potass. iodidi.....ʒ 4
Syr. sarsap. co.....
Aquæ.....aa fl. ℥ 2
Mix. Dose: a teaspoonful. *Dr. R. W. Taylor.*

In Syphilitic Cachexia.

℞ Iodoformi.....ʒ j.
Hydrarg. chlorid. corros.....gr. j.
Ferri redacti.....ʒ j.
M Ft. pil. No. xx. Sig.—One pill three times a day. *Bartholow.*

OR

℞ Iodoformi.....
Chinoidin.....
Ferri redacti.....aa ʒ j.
M. Ft. pil. No. xx. Sig.—One pill three times a day.

The above formulæ, more commonly than is supposed by advocates of special plans of treatment, exercise a most favorable influence over the course and duration of syphilis.

Bichloride of Mercury.

℞ Hydrarg. chlor. corros.....gr. 1
Potass. iodidi.....ʒ 2
Tr. gent. co.....fl. ℥ 4
Mix. Dose: a teaspoonful.

Iodide of Potash Mixture.

Charity Hospital, Blackwell's Island, N. Y.

℞ Potass. iodid.....ʒ 4
Syr. sarsap. co.....
Tinct. gent. co.....aa fl. ℥ 1
Mix. Dose: a teaspoonful.

Injections in Second Stage of Gonorrhœa—the Stage of Stasis.

℞ Zinci acetatis.....grs. 2–5
Aquæ destil..... ℥ j

OR

Acidi borac.....grs. 7–14
Aquæ destil..... ℥ j

Dr. F. R. Sturgis.

Inflammatory Stage of Gonorrhœa.

℞ Ol. cubeb.....♏ x–xv.

OR

Homeopathic tinct. cannabis sativa...♏ v–x.

This amount every two or three hours.

Dr. F. R. Sturgis.

In Gonorrhœa.

℞ Zinci sulphat.....
Acidi tannici.....ana gr. xv.
Aquæ rosæ..... ℥ vj.

Misce. Fiat injectio. To be used two or three times daily. *M. Ricord.*

In Gonorrhœa.

℞ Copaibæ..... ʒ ij.
Cubebæ..... ℥ ss.
Ceræ alb.....q. s.

Misce. Fiant pil. cxx. Ten pills three times a day, afterwards four times a day. *Niemeyer.*

In Gonorrhœa (repeated attacks).

℞ Cubebæ..... ℥ ss.
Copaibæ.....q. s.
Fiat pasta. A teaspoonful three times a day.

Mr. Erichsen.

To Prevent Chordee.

℞ Camphoræ.....gr. xv.
Opii pulv.....gr. iij.
Fiant pil. vj. Two or three to be taken every night at bedtime. *Ricord.*

In Syphilis (when Potas. Iodidum fails).

℞ Ammon. iodidi.....gr. viij.
Inf. aurantii..... ℥ j.
Misce. To be taken three times a day.

Mr. Berkeley Hill.

In Gonorrhœa, Acute Stage.

℞ Antim. tartarati.....gr. $\frac{1}{12}$
Magnes. sulph..... ʒ j.
Potass. nitrat.....gr. v.
Aquæ..... ℥ j.
Fiat haustus. To be repeated every fourth or sixth hour. *Mr. Erichsen.*

In Early Stage of Syphilis.

℞ Hydrarg. iodidi viridis.....gr. vj.
Confect. rosæ.....q. s.
Misce, et divide in pil. vj. One pill three times a day. *Mr. Erichsen.*

In Gonorrhœa.

℞ Tannin..... ʒ iss.
Vini burgundicæ.....Oiss.

Fiat injectio. One-third to be injected; if result unsatisfactory, the remainder. *Niemeyer.*

In Syphilis.

℞ Hydrarg. proto-iodidi.....
Lactucarii.....āā ʒ iss.
Ext. opii.....gr. ix.
Ext. guaiaci..... ʒ j.

Misce. Fiant pil. xxxvj. One pill twice a day.
Ricord.

Syphilitic Cachexia.

℞ Ext. sarsæ liquidi..... ℥ ss.
Acidi nitro-hydrochlor. dil.....♏ xx.
Aquæ..... ℥ iv.

Fiat haustus. To be taken three times a day.
Mr. Erichsen.

In Gonorrhœa.

℞ Liq. potassæ..... ʒ iiiss.
Potass. iodidi.....℈ j.
Pulv. rhei.....℈ ij.
Aquæ camphoræ..... ℥ vj.

Misce. Capiat ℥ j. ter die. *Mr. Milton.*

Syphilitic Coryza.

℞ Hydrarg. chlor. mit.....
Lycopodii.....aa ʒij

M. Sig.—Use as snuff three times daily, in syphilitic lesions of nose. *Gross.*

Mixed Treatment.

℞ Hydrarg. chlor. corros.....gr. j.
Potass. iodidi.....ʒij.
Tr. gentian. comp.....f℥iij.

M. Sig.—A teaspoonful three times a day.

Charity Hospital, N. Y.

Strangury.

℞ Decoct. uvæ ursi.....f℥viij.
Liq. potassæ. .....gtt. cxxx.
Tr. belladonnæ.....gtt. xlviij.

M. Sig.—Tablespoonful every four hours. *Agnew.*

Syphilis.

℞ Hydrarg. prot.....gr. vj.
Ft. pil. No. xxiv.

Sig.—One pill three times a day; every second day increase by one pill until first symptoms of ptyalism appear; then cut down dose one-half and continue for eighteen months this tonic dose; after that give—

℞ Potass. iodid.....ʒiss–iv.
Hydrarg. chlor. corros.....gr. i–iss.
Syr. aurant. cort.....f℥j.
Aquæ.....q. s. ad f℥iij.

M. Sig.—Teaspoonful three times a day for from six to twelve months. *Martin.*

Syphilis (Rebellious Cases).

℞ Acid. nitro-muriat. dil.....f ʒ iiss.
Syr. stillingiæ co..... ...f ʒ xiiiss.
Aquæ.....f ℥ ij.

M. Sig.—One or two teaspoonfuls three times a day in cases saturated with approved remedies, but still presenting mucous patches. *Bartholow.*

Mixed Treatment.

℞ Potassii iodidi.....gr. ij.
Syr. ferri iodidi.....♏ v.
Hydrargyri chloridi corros.....gr. $\frac{1}{64}$
Tr. nucis vom.....♏ ij.
Liq. arseni et hydrarg iodidi.....♏ ij.

M. Ft. pill or tablet.

Sig.—One, three times a day after meals with a half glass of water.

This is one of the best prescriptions for mixed treatment in solid form, the only form in which many patients can be induced to continue treatment for long periods. The tablet should be swallowed with a large draught of water. When well diluted the iodides are frequently tolerated by patients who could not otherwise take them.

Syphilitic Sore Throat.

℞ Hydrarg. chlorid. corros.....gr. iss.
Ac. muriat. dil.....ʒ j.
Glycerini.....℥ j.
Aquæ..... .....q. s. ad ℥ x.

M. Sig.—Use three or four times a day as a gargle.

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

VIII.

TONIC PREPARATIONS

The following tonic pills are much prescribed at the Gynæcological clinic of the Hospital of the University of Pennsylvania:

℞ Acid arseniosi..... } aa gr. $\frac{1}{30}$.
Strychniæ sulph..... ..... }
Ext. belladonnæ.....gr. $\frac{1}{6}$
Cinchonæ sulph..... .....gr. jss.
Pil. ferri carb.....gr. ijss.
M. Et ft. pil., No. j.

℞ Acid. arseniosi.....gr. $\frac{1}{30}$.
Cinchonæ sulph.....gr. jss.
Ferri et potass. tart.....gr. ij.
M. Et ft. pil., No. j.

The sulphate of cinchonia in these pills may be advantageously substituted by a proportionate dose of sulphate of quinia, the former being used simply on account of its cheapness. One pill may be given after each meal.

Tonic in Malarial Cachexia.

℞ Quiniæ sulphatis.....gr. ij.
Acidi arseniosi.....gr. $\frac{1}{30}$.
Pulv. capsici.....gr. j.
Ext. taraxaci.....q. s.
To be taken before each meal.

Where Anæmia is a Prominent Symptom in Malaria.

℞ Ferri et potassæ tart .....gr. v.
Liquoris potassii arsenitis .....m. ij.
Potassii bicarb.....gr. x.
Tr. nucis vomicæ.....♏ v.
Aquæ.....ad ʒ j.

To be taken in a wineglassful of water before eating. *J. O. Webster, M.D.*

Tonic.

℞ One rennet washed and chopped.....
Sherry wine.....O.j.
Macerate for twelve days, decant and filter and add
Acid. muriat. dilut.....
Tinct. nucis vomicæ.....aa ʒ ij.
Bismuthi subnit.....ʒ ij.

M. Sig.—One teaspoonful in a quarter of a tumbler of water before each meal. *Dr. T. G. Thomas.*

This is Dr. Thomas' formula for a digestive tonic in metritis, and its success commends it to all for the treatment of this most frequent malady.

Pil. Ferri. Quin. et Strych.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Quiniæ sulphat.....
Ferri redacti.....aa ʒ ı
Strychniæ acetat.....gr. ı
Ext. gentian.....q. s.

Mix. Divide into 60 pills.

Muriatic Acid Mixture.

Outdoor Department of Bellevue Hospital, N.Y.

℞ Acidi muriatici.....fl. ʒ 3
Tinct. gent. co.....
Aquæ.....aa fl. ℥ 8
Mix. Dose: a teaspoonful.

Ferruginous Aperient.

℞ Magnes. sulph.....℥ ij.
Ferri sulph.....gr. xxiv.
Acidi sulph. dil.....ʒ jj.
Infus. calumbæ.....ad ℥ viij.
Take two tablespoonfuls every night.

Dr. Farquharson.

Mist. Ferri Comp. (U. S. Pharm.)

(*Griffith's Mixture.*)

℞ Ferri sulphat.....gr. 20
Potassii carbon.....gr. 25
Pulv. myrrhæ.....
" sacchari.....aa ʒ 1
Spts. lavand. co.....fl. ʒ ½
Aquæ cinnamon. q. s. ad.....fl. ʒ 8
Mix. Dose: 1–2 tablespoonfuls. In preparing this, the sulphate of iron, dissolved in ½ fl. oz. of the cinnamon water should be added last. The mixture should be made fresh when wanted.

"Smith's Bitters."

℞ Tinct cinch. co.....
Tinct. gent. co.....aa fl. ℥ 1
Mix. Dose: a teaspoonful.

Wine of Iron and Citrate of Quinidiæ.

℞ Ferri et quinidiæ citr..... ʒ i
Vini xerici.....fl. ℥ ii

Mix. Dose: a teaspoonful. To prevent confusion this should always be prescribed by writing the word "Quinidiæ" in full.

Strychnia Mixture.

(*Hall's Modified.*)

℞ Strychniæ acet.....gr. i
Tr. cardam. co.....fl. ʒ ½
Alcoholis.....
Aquæ.....aa fl. ʒ ii½
Syrupi q. s. ad.....fl. ℥ iv

M. Dose: a teaspoonful.

The following tonic formulas are very highly recommended by Dr. Wm. Goodell:

Basham's iron mixture, with the addition of fractional doses of strychnia, will be found very admirable in its effects. There are so many indifferent recipes for making this celebrated mixture that I shall here give the one which seems to me to be the best:

℞ Tinct. ferri chloridi.....fl. ʒ iij.
Acid. acetic. diluti.....fl. ℥ ss.
Liquor ammoniæ acetat.. .....fl. ℥ iijss.
Curacoæ.. .....
Syrupi simplicis.....aa fl. ℥ j.
Aquam. ad.....fl. ℥ viij

M. Sig.—One tablespoonful after each meal.

The following formula makes another very elegant and generally useful preparation of iron:

℞ Tinct. ferri chloridi.....fl. ʒ ij.
Acid. phosphorici diluti.....fl. ʒ iij.
Spts. limonis.....fl. ʒ j.
Syrupi simplicis.....fl. ℥ ijss.
Aquam, ad.....fl. ℥ vj.
M. Sig.—One tablespoonful after each meal.

The dilute phosphoric acid is added both because it is a valuable nerve-tonic and because it has the property of disguising the styptic taste of the iron; so much so, that children readily take this mixture.

There are two other tonic preparations which we prescribe very frequently in the Hospital of the University of Pennsylvania, and with capital results. One of them is Blaud's pill, which Niemeyer extols so very highly.

℞ Pulv. ferri sulphat. exsiccat.....
Potass. carb. puræ.....aa ʒ ij.
Syrupi.....q. s.
Ut fiat massa dividenda in pilulas, No. xlviij.

During the first three days one pill is to be taken after each meal. On the fourth day four pills are taken during the day, on the fifth day five pills, on the sixth day six; that is to say, two pills after each meal. For three days more six pills are taken daily; then the dose is to be increased by one pill daily until three pills are taken after each meal. On this final dose the patient is kept for three or four weeks, as the case may be. In stubborn cases I have occasionally run up the dose to the number of five pills thrice daily, and have seen no other bad effects from it than

a feeling of fulness in the head. This immunity is probably owing to the conversion of the iron sulphate into a carbonate.

The other preparation is a valuable alterative tonic, for the formula of which I am indebted to my friend Dr. A. H. Smith.

℞ Hydrarg. chloridi corrosivi.....gr. i.—ij.
Liq. arsenici chloridi.....fl. ℥ j.
Tinct. ferri chloridi.....
Acid. hydrochloridi dil.....aa fl. ʒ iv.
Syrupi.....fl. ℥ iij.
Aquam, ad.....fl. ℥ vj.

M. Sig.—One dessertspoonful in a wineglassful of water after each meal.

Anæmic and chlorotic patients will fatten and thrive wonderfully on this mixture. I call it the Mixture of Four Chlorides. It should not be given for a longer period than two weeks at a time.

Wine of Iron and Citrate of Quinine.

℞ Ferri et quiniæ cit.....ʒ i
Vini xerici.....fl. ℥ 2

Mix. Dose: a teaspoonful.

Iron and Citrate of Ammonia.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Ferri et ammon. cit.....
Ammonii carbon.....aa gr. 32
Syrupi.....
Aquæ anisi.....aa fl. ℥ 2

Mix. Dose: a teaspoonful. *Dr. J. L. Smith.*

Lemon Tonic.

Charity Hospital, Blackwell's Island, N. Y.

℞ Cinchonæ sulph.....gr. 30
Acidi sulph. dil.....q. s.
Aquæ.....fl. ℥ 1
Acidi citrici..... ʒ ½
Syrupi.....fl. ℥ 1½
Tinct. ferri chlor.....fl. ʒ ½
Aquæ q. s. ad.....fl. ℥ 4
M. Dose: a teaspoonful.

Iron and Cinchona.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Cinchonæ sulph..... ʒ 1
Tinct. ferri chloridi.....fl. ʒ 2
Aquæ q. s. ad.....fl. ℥ 4
Mix. Dose: a teaspoonful.

Sulphate of Cinchona.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Cinchonæ sulph..... ʒ 1
Acidi sulphur. dil.....q. s.
Aquæ.....fl. ℥ 4
Mix. Dose: a teaspoonful.

Iodide Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Potassii iodidi..... ʒ 4
Syr. ferri iodidi.....fl. ℥ 1
Tinct. calumbæ q. s. ad.....fl. ℥ 4
Mix. Dose: a teaspoonful.

Loomis' Tonic.

℞ Quiniæ sulphat.....gr. 30
Acidi sulph. dil.....q. s.
Aquæ.....fl. ℥ 2
Tinct. ferri chlor.....fl ℥ ½
Spts. chloroformi.....fl. ʒ 6
Glycerinæ q. s. ad.....fl. ℥ 4
Dose: a teaspoonful.

Elixir of Cinchona and Iron.

℞ Extr. cinchonæ fl.....fl. ℥ 1
Ferri et ammonii citr..... ʒ 2
Spiritus aurantii (1:10).....fl. ℥ 1
Alcoholis.....fl. ℥ 4
Ol. cinnamoni.....m. 15
Aquæ.....fl. ℥ 6
Syrupi q. s. ad.....fl. ℥ 16

With the oil of cinnamon and water make water of cinnamon; dissolve in this the ammonio-citrate of iron; add the spir. of orange and the fl. ext. of cinchona. Filter, and add enough syrup to make the product measure one pint. Dose: a teaspoonful.

Iron and Quinine.

℞ Quiniæ sulph.....gr. 30
Acidi sulph. dil.....q. s.
Aquæ.....fl. ℥ 2
Tinct. ferri chlor.....fl. ʒ 2
Dose: a teaspoonful.

Wine of Citrate of Iron.

℞ Ferri et ammonii citratis.....gr. 20
Vini xerici.....fl. ℥ 2
Mix. Dose: a teaspoonful.

Thomson's Tonic.

Bellevue Hospital, New York.

℞ Ferri et ammon. cit..... ʒ 1
Ammon. carbonat.....gr. 30
Tr. gentian co.....
Tr. quassiæ.....aa fl. ℥ 2
Syrupi.....fl. ℥ 1½
Aquæ q. s. ad.....fl. ℥ 8
Mix. Dose: a dessertspoonful.

Dr. W. H. Thomson.

Strychnine Tonic.

Insane Asylum, Blackwell's Island, N. Y.

℞ Tinct. ferri chlor.....
Tinct. nucis vom.... .....aa m. 10
Aquæ q. s. ad.....fl. ʒ 1
Mix. One dose. To be taken thrice daily, after meals.

Tonic and Stimulant.

℞ Liq. potass. arsenitis.....ʒj.
Vini ferri amar.....℥vj.

M. Sig.—Tablespoonful three times a day after meals.

F. P. Henry.

For Anaemia.

℞ Mangani dioxidi.....gr. j.
Ferri reducti.....gr. j.
Ext. gentianæ.....gr. j.
Ext. cascaræ sagrad.....gr. ss.
Ext. nucis vom.....gr. ⅛
Capsici.....gr. ⅛
M. Ft. pil.

Sig —One three times a day after meals. An excellent hematic tonic for women and young girls, particularly when accompanied by scanty or painful menstruation.

Blaud's Ferruginous Pills.

℞ Ferri sulphatis exs.....
Potassii carb.....aa gr. iiss.
M. Ft. pil.
Sig.—One or two after meals.

These pills have justly come to be regarded as one of the most efficient, if not the most efficient agent in the treatment of *chlorosis* and the anæmias of women and young girls. They were first brought to the notice of the profession by Niemeyer, who thus wrote of them: "For more than twenty years I have used these pills almost exclusively in chlorosis, and have witnessed such brilliant results from them in a large number of cases, that I have never needed any opportunity to experiment with other articles. At Magdeburg and Greifswald I often had to send my receipt for the pills to a great distance, my good fortune in the treatment of chlorosis—to which, by the by, I owe the rapid growth of my practice—having given me a great repute as the possessor of a sovereign remedy against this disease."

The activity of Blaud's pills is increased by combination with other tonics. The following combination proposed by Hare is one of the most powerful alterative and reconstructive tonics at our command. It is now made up into tablet triturates by several leading firms in full strength and half strength.

℞ Mas. ferri carb. (Blaud).....gr. v.
Hydrarg. chloridi corros.....gr. $\frac{1}{80}$
Strychninæ sulphatis .....gr. $\frac{1}{60}$
Ac. arseniosi.....gr. $\frac{1}{60}$
Capsici .....gr. $\frac{1}{64}$
Ext. gentiani .....gr. $\frac{1}{16}$
M. Ft. pil. No. i.
Sig.—One after meals.

The following combination is designed to overcome the constipating effects observed in some patients.

℞ Mas. ferri carb. (Blaud).....gr. iii.
Quininæ sulphatis.....gr. ss.
Ext. nucis vomicæ.....gr. $\frac{1}{10}$
Aloin.....gr. $\frac{1}{10}$
M. Ft. pil. No. i.
Sig.—One or two three times a day after meals.

Tonic for Malaria.

℞ Ferri reduct.....gr. ij.
Quininæ sulphat.....gr. ij.
Strychniæ sulphat.....gr. $\frac{1}{50}$
Ac. arseniosi.....gr. $\frac{1}{50}$
Pulv. aloes.....gr. $\frac{1}{10}$
M. Ft. pil. Sig.—One every three to four hours for chronic malaria.

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

IX.

EXTERNAL APPLICATIONS.

For Fissured Nipples.

℞ Liq. ferri subsulphatis..... ʒ ij.
Glycerini..... ʒ vj.
M. Sig.—Apply with a camel's hair brush to affected part. *Bartholow.*

In Parasitic Skin Diseases.

℞ Sodii hyposulphitis..... ..... ℥ iij.
Acid. sulphurosi dil..... ℥ ss.
Aquæ q. s. ad..... ℥ xvj.
M. Sig.—As a lotion. *Dr. Startin.*

OR

℞ Potass. sulphureti..... ..... ℥ ss.
Aquæ calcis..... ℥ xvj.
M. Sig.—As a lotion. *Bartholow.*

Chilblains.

℞ Acid. sulphurosi..... ʒ iij.
Glycerini..... ʒ j.
Aquæ..... ..... ʒ j.
M. Sig.—As a lotion. *Bartholow.*

Pruritus.

There are several anti-pruritic powders, which are sometimes indispensable. The most important is that which is commonly called Anderson's powder. It is formed as follows:

℞ Pulv. amyli..... ℥ j.
Pulv. camph..... ʒ jss.
Zinci oxid..... ℥ ss.

This must be carefully made, the camphor being reduced to an impalpable powder and then thoroughly incorporated with the other ingredients. It may be either lightly dusted upon the parts or it may be quite copiously rubbed into the meshes of linen lint and then applied. The proportions may be altered, either by increase or diminution.

Dr. R. W. Taylor.

Ung. Hydr. Ox. Rub. c. Plumbo.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Hydrarg. oxidi rubri.....
Plumbi acetat.....aa gr. 8
Cerati..... ℥ i

Mix. *Dr. McKay.*

Sulphur Paste.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Sulphuris sublimati..... ℥ i
Ætheris.....fl. ʒ ʒ
Glycerinæ.. .....fl. ℥ i

Mix.

Ung. Hydrarg. Oxid. Rubri.

℞ Hydrarg. oxid. rubri.....gr. 60
Oleo-paraffini (vaseline)..... ℥ 1
Mix.

In Parasitic Skin Diseases.

℞ Hydrarg. chlor. cor.....gr. iv.
Alcohol ..... ʒ vj.
Ammon. muriat..... ʒ ss.
Aquæ rosæ q. s. ad..... ℥ vj.
M. Sig.—As a lotion. *Tilbury Fox.*

Glue Burn Mixture.

℞ White glue..... ℥ 7½
Water, cold .....O.j.
Glycerin.....fl. ℥ 1
Carbolic acid.....fl. ʒ 2

Soak the glue until it is soft; then heat on water bath until melted; add the glycerin and carbolic acid, and continue heating until in the intervals of stirring a glossy strong skin begins to form over the surface. When wanted for use, heat on water bath until melted, and apply with a flat brush over the burned part.

Ung. Picis Alkalinum.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Liquoris picis alkalini.....fl. ʒ 1
Cerati..... ℥ 1
Mix.

Goa Ointment.

℞ Pulv. arariobœ (Goa powder)...gr. 50 to 200
Oleo-paraffini (vaseline).....gr. 500

Melt the vaseline on a water bath, add the Goa powder, stir and heat for about ten minutes; then strain quickly into a capsule standing on ice, and stir briskly until cold.

In the Pruritus of Pregnancy.

℞ Thymol.....gr. xv.
Vaseline.. .....gr. xxx.
Powdered brick clay..... ℥ iij.

Dissolve the thymol in the vaseline and rub it up with the clay. *Prof. Montrose A. Pallen, M.D.*

This is to be applied to the pruritic parts, washed off every day or two and re-applied.

Dr. Pallen's experience has been, that excepting those cases depending on trophic nerve causes, this prescription will always effect a cure. He advises its use also in herpes and similar eruptions accompanying the later months of gestation.

Bleeding Hemorrhoids.

℞ Pulv. aluminis..... ʒ ij.
Pulv. camphoræ.....
Pulv. opii.....aa ʒ j
Unguent..... ..... ℥ j

M. Sig.—Ointment. *Bartholow.*

When the hemorrhoids protrude, bleed, and are painful, the above will be found a soothing and astringent application.

Emplastrum Capsicum.

(*McCready's Plaster.*)

Spread adhesive plaster with oleo resina capsici, leaving, however, a narrow margin all around free.

Dr. B. W. McCready.

Lotion for Iritis.

℞ Morphiæ sulph.....gr. iv.
Zinci sulph.....gr. iii.
Atropiæ sulph.....gr. ii.
Aquæ destillat..... ℥ j.
M. Sig.—As a lotion. *Dr. Bartholow.*

Sore Nipples.

℞ Aquæ rosæ.....
Glycerine.....aa ℥ ij.
Acidi tannici..... ʒ ij.
Ft. lotion.
Sig.—Soak lint in this solution and apply to nipples.

Dr. Barker.

If the ulcerative process has commenced, it is advisable to stop nursing and paint the nipple with a solution of nitrate of silver, 10 gr. to the ℥ of distilled water.

As a Hæmostatic.

℞ Acidi tannici.....℈ j.
Aluminis.....℈ ij.
Aq. rosæ..... ℥ iij.
M. Sig.—For external use as a hæmostatic.

Dr. Monsel.

Pruritus Vulvæ.

℞ Sodii hyposulphitis..... ʒ iv.
Glycerini..... ʒ ij.
Aquæ destillat.....ad ℥ vj.
M. Sig.—As a lotion. *Fox.*

This simple combination has proved very effective in that troublesome and annoying malady, pruritus vulvæ and also in tinea versicolor.

In Amenorrhœa from Anæmia and in Chlorosis.

℞ Pulv. ferri sulph.....
Potass. carb. puræ.....aa ʒ ij.
Mucil. tragacan..... .....q. s.
M. Div. in pil. No. 48.

Sig.—To be given daily in doses gradually increasing until three pills are taken after each meal.

To counteract the costive effect of the sulphate of iron in this, we may give:

℞ Pulv. glycerrhizæ rad.....
Pulv. sennæ.....aa ℥ ss.
Sulph. sublim.....
Pulv. fœniculi.....aa ʒ ij.
Sacchar. purif..... ℥ jss.

Sig.—Teaspoonful in half a cupful of water at bedtime.

Where the disease is due to torpidity of the ovaries, the following may be used with advantage:

℞ Ext. aloes..... ʒ j.
Ferri sulph. exsiccat..... ʒ ij.
Assafœtida..... ʒ iv.

Sig.—One pill after each meal, this number to be

increased gradually to two or three after each meal. If bowels are over-active, return to the initial dose of one pill.

The above formulæ are those used in the Hospital of the University of Pennsylvania, Philadelphia, and have been proven by thorough trial to be most efficient.

In Inflammation of the Nipples.

℞ Emplastri plumbi..... ʒ ij.
Ætheris sulphurici..... ʒ ss.
Collodion flexile..... ℥ j.

M. Sig.—Powder the lead plaster, add the ether and mix them well together before adding the collodion. It makes a creamy mixture, and is to be applied with a brush over every portion of the carefully dried nipples, with the exception of the openings of the milk ducts. *Dr. Albert H. Smith.*

This formula is highly commended by Dr. Smith as a protective application.

Dr. R. W. Taylor's Lotion.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Sulphuris sublimati..... ʒ ʒ
Spts. camphoræ.....fl. ʒ ʒ
Sodii biboratis..... ʒ ʒ
Glycerinæ..... .....fl. ʒ ʒ
Aquæ..... ℥ ʒ

Mix.

Conjunctivitis.

℞ Zinci sulphat.....gr. ij.–viij.
Morphiæ sulphat.....gr. ij.–iv.
Atropiæ sulphat.....gr. ss.–j.
Aquæ rosæ..... ℥ j.
M. Sig.—For the eye. *Dr. Bartholow.*

To Destroy the Odor of Foul Breath, the Smell of the Axilla, and the Fetor of the Sweat of the Feet.

℞ Potass. permangan.....gr. vj.
Aquæ..... ʒ vj.
Sig.—Apply frequently.

It is a fact too little appreciated by physicians that success in practice often depends more on attending to some such trivial affection as the above, than on the successful management of a complicated medical or surgical case.

Carson's Paint.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Olei tiglii.....fl. ʒ ½
Ætheris.....fl. ʒ i
Tinct. iodinii co.....fl. ʒ 2½
M. Counter irritant and vesicant in pleurisy, etc.

Local Application for Chilblains.

℞ Acid carbol..... ʒ i.
Tinct. iodinii..... ʒ ij.
Acid tannici..... ʒ ii.
Cerat. simplicis..... ℥ iv.
M. Sig.—Ointment. *Dr. Bartholow.*

Eczema.

℞ Liq. plumb. subacetat..... ℥ j.
Glycerini..... ℥ ss.
Cherry laurel water..... ℥ iijss.
Sig.—Lotion. *Dr. Bartholow.*

This will be found to be very serviceable in eczema, characterized by great heat and redness and excessive discharge.

Facial Erysipelas.

℞ Quiniæ sulph..... ʒ ss.
Belladonna ext.....gr. iij.
M. Ft. pil. No. x.
Sig.—One every six hours. *Dr. Bartholow.*

Croton Oil Liniment.

Ninety-ninth Street Reception Hospital, New York.

℞ Olei tiglii.....fl. ʒ 2
Olei olivæ.....
Olei terebinthinæ.....
Aquæ ammoniæ.....
Spiritus camphoræ aa p. e. q. s. ad.....fl. ℥ 2
Mix. Sig.—Externally in chronic muscular pains.

Ung. Diachylon Hebræ.

℞ Emplast. plumbi..... ℥ 5
Olei olivæ.....fl. ℥ 4
" lavandulæ.....fl ʒ 1

Melt the lead-plaster and olive oil together at a gentle heat; then stir until the mixture begins to stiffen, and incorporate with it the oil of lavender.

Ung. Acidi Chrysophanici.

℞ Acidi chrysophanici.....gr. 20
Oleo-paraffini (vaseline).....gr. 190

Melt the vaseline on a water-bath, add the acid, stir and heat for about ten minutes; then strain quickly through muslin into a capsule standing on ice, and stir briskly until cold.

Buck's Burn Mixture.

℞ Pulv. tragacanthæ..... ℥ 2
Pulv. acaciæ..... ℥ 4
Syrupi fusci.....
Aquæ bullientis.....aa O1

Mix.

Lotion for Syphilitic Eruptions.

℞ Hydrarg. chlor. cor.....gr. iv.
Acid. nitric. dil..... ʒ j.
Acid. hydrocyanici dil..... ʒ j.
Glycerini..... ʒ ij.
Aquæ..... ℥ viij.

M. Sig.—Use as a lotion. *Dr. Startin.*

Carron Oil.

(Lime Liniment. Linimentum Calcis, U. S. Ph.)

℞ Olei lini..... ℥ 7
Aquæ calcis.....fl. ℥ 8

Mix.

Lotion in Acute Eczema.

℞ Potass. bicarbonat..... ʒ ss.
Aquæ.....Oj.
Use as a lotion. *Farquharson.*

As a Depilatory.

℞ Quicklime..... ℥ ss.
Yellow sulphide of arsenic.....gr. xx.
Starch.....gr. clxxx.
M. Sig.—Apply as a paste. *Bartholow.*

A preparation of this kind is probably such a one as that used by the Egyptian women to remove the hair from their pubes.

Lotion in Acne.

℞ Hydrarg. chlor. cor..... ℈ j.
Glycerini..... ℥ ss.
Spts. vini. rect..... ℥ viij.
Spts. rosemar..... ʒ iv.
M. Sig.—Use as a lotion. *Bartholow.*

Acne Indurata.

℞ Hydrarg. iod. rubri.....gr. v.
Adipis..... ℥ j.
M. Sig.—As an ointment. *Bartholow.*

For Sore Nipples.

℞ Plumbi nitrat.....gr. x–xx.
Glycerine..... ℥ j
Sig.—Rub on the affected part.
Dr. Fordyce Barker.

℞ Aquæ rosæ..... ℥ iijss.
Liq. plumb. subacetat dil..... ℥ ss.
Ext. opii. aq..... ʒ j.

Sig.—Use as a lotion, wetting linen in this and covering the breast with two thicknesses of it, having previously applied a bread and milk poultice to the breast.

This is recommended by Dr. Barker in inflamed nipples.

For Freckles, Sunburn, and Tan.

℞ Potass. carbonat..... ʒ iij.
Sodii chloridi..... .. ʒ ij.
Aquæ rosæ..... ℥ viij.
Aquæ aurantii flor..... ℥ ij.
M. Sig.—Lotion. *Bartholow.*

In Itch, Impetigo, etc.

℞ Creasoti..... ʒ ss.
Aquæ destil.....Oj.
Fiat lotio. *Dr. Dunglison.*

To Prevent and Cure Chilblains.

℞ Cupri sulphatis..... ℈ ss.
Aquæ destil.....f ℥ j.
Fiat lotio.

To be applied with a camel-hair pencil to the part affected; when dry, smear the part with spermaceti ointment. Repeat the application for two or three days, till the skin appears to be affected.

Dr. Graves.

In Sloughing Phagedæna.

℞ Cupri sulph.....gr. j.
Ext. opii.....gr. v.
Aquæ..... ℥ j.
Fiat lotio. *Mr. Erichsen.*

In Impetigo or Acne.

℞ Acidi carbolici.....gr. x.
Glycerini.....
Aquæ rosæ.....āā ℥ j.
Fiat lotio. *Dr. Headland.*

In Ringworm.

℞ Thymol..... ʒ j.
Adip. præp..... ℥ j.
Ft. unguentum. *Mr. Balmanno Squire.*

In Pruritus Ani.

℞ Acidi carbolici.....gr. vj.
Aquæ..... ℥ j.
Fiat lotio. *Mr. Christopher Heath.*

In Scabies.

℞ Sulph præcipitati.....gr. xxx.
Ol. santal. flav.....♏ ij.
Adipis præparati..... ℥ j.
Fiat unguentum. *D. R. Liveing.*

For Bedsores.

℞ Hydrarg. perchlor.....gr. ij.
Sp. rectificat..... ℥ j.
Fiat lotio. *Mr. Erichsen.*

For Sore Nipples

℞ Hydrarg. subchlor..... .....gr. xxx.
Magnesiæ carb..... ℈ j.
Ung. rosæ..... ℥ j.
Fiat unguentum. *Dr. Fordyce Barker*

In External Piles.

℞ Ung. zinci..... ℥ j.
Liq. plumbi subacet.....
Liq. opii sedat.....ana ʒ ss.
Misce. Fiat unguentum, bis in die parti affectæ applicandum. *Mr. Coulson.*

In Prurigo Senilis.

℞ Ol. staphisagriæ..... ʒ j.
Adipis præp..... ℥ j.
Fiat unguentum. *Mr. Balmanno Squire.*

In Obstinate Acne.

℞ Hydrarg. oxidi rubri.....
Hydrarg. ammoniati.....ana gr. v.
Adipis præp..... ℥ j.
Fiat unguentum. *Dr. Tilbury Fox.*

In Ringworm.

℞ Thymol..... ..... ʒ ss.
Etheris..... ..... ʒ v.
Sp. rectif..... ʒ iiss.
M. To be used as a wash every morning, and followed by an application of glycerine and perchloride of mercury. *Mr. Malcolm Morris.*

Anæsthetic Application.

℞ Aconitiæ.....gr. j.
Sp. rectif..... ʒ j.
Aquæ destil..... ʒ ix.

Solve aconitiam in spiritu, dein adde aquam, et cola.

Adde, glycerini..... ʒ ij.
Olei bergamotæ.....♏ v.

M. Fiat lotio.

The solution in spirit and water forms *Liquor Aconitiæ*, which may be prescribed internally in minute doses in acute rheumatism. With the glycerine, etc., a lotion is formed, which will rapidly produce numbness if rubbed on the painful part in a quantity of about ʒ ss. at a time. *Dr. Headland.*

Sensitive Ulcers of the Rectum.

℞ Chloroform.....f ʒ j.
Zinci oxidi..... ʒ ss.
Ol. olivæ..... ʒ j.
Cerati cetacei..... ʒ iv.

Misce. Fiat unguentum. *Mr. Curling.*

In Eczema, Herpes, and Simple Excoriations.

℞ Pulv. sabinæ.....
Pulv. æruginis, ana partes æquales.....

Fiat pulvis. To be dusted over the part affected.

Mr. Acton.

In Acne Between Menstrual Periods.

℞ Sulphuris..... ʒ j.
Glycerini..... ℥ j.
Aquæ rosæ..... ℥ x.
Misce. Fiat lotio. *Dr. Ringer.*

To Prevent Pitting in Vaccinia.

It has been recommended to touch each pustule with nitrate of silver or camphor, or to bathe the face with a solution of four scruples of nitrate of silver to ℥ j. water. A mercurial plaster formed of

Ung. hydrarg.....25 parts.
Yellow wax.....10 "
Black pitch..... 6 "

has a good effect. Dr. Aitken gives this as the formula used at the Children's Hospital in Paris. According to Guersant, four or five days are sufficient for the application to prevent pitting.

Carron oil is a good application till the scabs begin to loosen; they should always be removed when dry, or they stain the skin permanently. Mr. Marson recommends cold cream and oxide of zinc, or if the discharge be thin and excoriating, calamine mixed with olive oil. Dr. W. Stokes uses light poultices over the face or a mask of lint soaked in glycerine and water and covered with a further mask of oiled silk.

E. Ellis.

Pruritus.

℞ Chloral hydrat.....
Pulv. camphoræ..... .....aa ʒiv.

M. Sig.—Apply with a camel's-hair brush for itching or local pain. *George Bird.*

Eczema of the Hands.

℞ Acid. salicylic.....gr. xxv.
Pulv. amyli.....
Pulv. zinci ox.....aa ʒij.
Petrolati..... .....℥ss.
M. Sig.—Use twice a day. *Stelwagon.*

OR

℞ Resorcin..... ...gr. xl.
Glycerinæ .....♏xv.
Alcohol..... ... .ʒj.
Aquæ.... ..... .ʒiv.
M. Sig.—To be used in conjunction with an ointment. *Stelwagon.*

Eczema.

℞ Ol. cadini.....℥ss.
Glycerinæ.....ʒj.
Ungt. diachyli.....℥iiss.
M. Sig.—Apply locally in squamous eczema with t' ickened skin. *Tilbury Fox.*

Tinea Versicolor.

℞ Acid. salicylici.....gr. xxx.
Sulphuris loti.....ʒiiss.
Lanolini..... ..ʒxxv.
M. Sig.—Apply with friction. *Liebreich.*

OR

℞ Hydrarg. chlor. corros.....℈j.
Saponis viridis.....℥ij.
Alcoholis.....℥iv.
Ol. lavandulæ. .....ʒj.
M. Sig.—To be rubbed in well night and morning. *Van Harlingen.*

Acne Rosacea.

℞ Ichthyol.....
Resorcin .....aa gr. xv.–lxxv.
Lanolin.....ʒvj.
Aq. dest llat.....q s. ad ℥iss.
M. Sig.—Apply topically.

Impetigo Contagiosa.

℞ Acid. salicylic.....gr. xxx.
Petrolati..... .....ʒj.
Zinci oxidi.....
Amyli.....aa ℥ss.
M. Sig.—Apply after removal of crusts and cleansing.

Lassar.

Urticaria-Pruritus.

℞ Chloralis..... .....
Camphoræ.....aa ʒj.
Pulv. amyli.....ʒi–ij.
M. Sig.—Keep tightly corked in a wide-mouthed bottle. Rub in with hand. *Bulkley.*

Erysipelas.

℞ Aristol.....gr. xx.
Collodii..... .....ʒj.
M. Sig.—Apply freely with a camel's-hair brush over and slightly beyond the inflamed area.

OR

℞ Ichthyol.....ʒj.
Lanolini.....ʒix.
M. Sig.—Apply freely over the inflamed area.

Nussbaum.

Alkaline Bath.

℞ Potass. carbonat..... ..... ℥ii–vj.
Sodii borat..... ℥ij.
Aq. ferventis (90°–95° F.).....cong. xx–xxx.

M. Sig.—Alkaline bath to be used in pruritus and various skin disorders. *Tilbury Fox.*

Frostbite.

℞ Ichthyol..... .....
Resorcin.....
Acid. tannic.....aa ʒj.
Aquæ..... ..... ʒv.

M. Sig.—Apply with a brush at night. *Boeck.*

Chilblains—Frostbite.

℞ Iodi..... .....℈j.
Potass. iodid.....gr. iv.
Aq. destillat.....♏vj.
Adipis..... .....℥j.

M. Sig.—Apply once daily. *Hebra.*

Contusions.

℞ Tr. capsici.....
Tr. myrrh..... .....
Tr. opium.....aa ʒij.
Tr. guaiac.....ʒj.
Spts. camphor.....℥ij.

M. Sig.—Use locally.

Freckles—Sunburn.

℞ Plumbi acetat.....gr. xv.
Acid. hydrocyanic. dil.....♏xx.
Alcoholis.....℥ss.
Aquæ.....q. s. ad ℥vj.
M. Sig.—Apply with a sponge. *Tilbury Fox.*

Dressing for Wounds.

℞ Iodoform.....ʒj.
Collodii flex.....ʒvij.
M. Sig.—Stitch the edges of the wound together and apply with a brush. *Bruns.*

Loss of Hair after Fever.

℞ Ex. jaborandi fl.....
Tr. cantharidis.....aa f ℥ss.
Glycerinæ.....
Ol. vaselini.....aa f ℥j.
M. Sig.—Apply thoroughly to the scalp at night. *Bartholow.*

For Odor and Sweating of Feet.

℞ Acid. salicylici.....gr. xlv.
Pulv. amyli.....ʒv.
Pulv. talc.....ʒxxij.
M. Sig.—Dust over the feet. (Used in the German army.)

To Prevent Bed-Sores.

℞ Alumin.....
Sodii chloridi.....aa ℥ss.
Aquæ.....
Alcoholis.....aa Oj.
M. Sig—For local use, twice daily. *Forbes.*

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

X.

COD LIVER OIL PREPARATIONS.

Cod Liver Oil Emulsion.

℞ Olei morrhuæ.....partes 28
Glyconini*.....partes 9
Spts. ammon. arom.....partes 1
Vini xerici.....partes 20
Spts. amygdal. amar.....partes 2

Mix. All to be taken by weight. Dose: a tablespoonful.

Phosphorated Emulsion of Cod Liver Oil.

℞ Olei morrhuæ.....partes 20
Olei phosphorati (1 per cent).....partes 2
Glyconini.....partes 7
Spt. ammon. arom.....partes 1
Syrupi.....partes 10
Acidi phosphor. dil.....partes 4
Spts. amygdal. amar.....partes 2

---

* *Glyconinum, or Glyceritum Vitelli.*

℞ Vitellorum ovorum.....partes 4
Glycerinæ.....partes 5

Beat or whip the yolks of the eggs, which must be fresh, in the usual manner, pour the liquid into a bottle, add the glycerine, and shake them well together. One pint of cod liver oil requires about 4 fl. oz. of glyconin to emulsionize it.

All by weight. Put the glyconin into a mortar and add the oils to it in very small quantity at a time, triturating the mixture actively and constantly. Then add the other ingredients in the order in which they are named. Dose: a tablespoonful.

Phosphorated Cod Liver Oil.

Bellevue Hospital, N. Y.

℞ Olei phosphorati * (1 per cent)..... gr. 100
Ætheris.....fl. ʒ 2
Olei morrhuæ q. s. ad.....fl. ℥ 16

Mix. Two hundred and thirty-three minims, or practically ½ fl. ℥, contain $\frac{1}{30}$ grain of phosphorus. The phosphorated oil should be weighed, not measured.

---

* *Oleum Phosphoratum.*

℞ Phosphori.....gr. 1
Olei morrhuæ.....gr. 99

This is a 1 per cent solution of phosphorus in cod liver oil, proposed by Dr. E. R. Squibb. It is made with the utmost care, and contains the full amount of phosphorus. If only a portion of the contents is to be used, a few drops of ether should be poured into the vial, before it is again corked and sealed. If a fine film should form on the surface or at the bottom, the oil must be poured out, so as to leave this in the vial. Should this film increase, or much of a precipitate make its appearance, a fresh bottle should be used.

It is best to add the whole contents of a bottle at once to sufficient cod liver oil to be ready for administration. The latter may be kept on hand in full and well closed bottles, which are to be kept in the dark.

When using the phosphorated oil it should always be taken by *weight.*

Agreeable Form of Taking Cod Liver Oil, for Children.

℞ Olei morrhuæ..... ℥ ss.
Mucilag. acaciæ..... ℥ ij.
Sacchari..... ʒ ij.
Tinct. lavand. comp..... ♏ xx.
Aquæ..... ℥ ss.
Dose: a teaspoonful. *Farquharson.*

Emulsion of Cod Liver Oil with Lime.

℞ Olei morrhuæ.....fl. ℥ 2
Aquæ calcis.....fl. ℥ 1½
Syr. calcis lactophosph.....fl. ℥ 1½
Mix. Dose: a teaspoonful. *Dr. Bosley.*

Cod Liver Oil Mixture.

Outdoor Department of Bellevue Hospital, N. Y.

℞ Olei morrhuæ.....fl ℥ 16
Liquor potassæ.....fl. ʒ 2
Mellis.....fl. ℥ 3
Pulv. acaciæ..... ℥ 1
Ol. anisi.....gtt. 20
Ol. menthæ vir.....gtt. 18
Mix. Dose: a tablespoonful. *Dr. Winston.*

Cod Liver Oil Emulsion.

Charity Hospital, Blackwell's Island, N. Y.

℞ Olei morrhuæ..... ...
Aquæ calcis.....aa fl. ℥ 8
Olei cinnamomi.....gtt. 10
Mix. Dose: a tablespoonful.

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

XI.

MISCELLANEOUS PRESCRIPTIONS.

Solution Ergotini.

Charity Hospital, Blackwell's Island, N. Y.

(Ergotin solution for hypodermic use.)

℞ Ergotini.....gr. 36
Glycerinæ.....
Aquæ .....aa m. 108
Mix.

Liquor Chloroform Co.

(B. Squire's Formula for "Chlorodyne.")

℞ Chloroformi purif.....fl. ℥ 4
Ætheris fort.....fl. ℥ 1
Alcoholis fort.....fl. ℥ 4
Syrupi fusci.....fl. ℥ 4
Extr. glycyrrh. pulv..... ℥ 2½
Morphiæ hydrochlorat.....gr. 8
Ol. menthæ piper .....m. 16
Acid hydrocyan. dil (2 per cent).....fl. ℥ 2
Syrupi.....fl. ℥ 17½

Dissolve the morphia and oil of peppermint in the alcohol; mix the chloroform and ether with this solution. Mix the liquorice with the syrup and add the molasses. Shake these mixtures well together, lastly

add the hydrocyanic acid, and again shake well. Dose: 10 to 20 drops. Always shake the mixture before using.

Formulæ for Hypodermic Administration of Quinine.

℞ Quiniæ sulphat..... ʒ j.
Morphiæ sulphat.....gr. ss.
Acid. sulphur. dil..... .....m. xl.
Aquæ destillatæ..... ℥ j.
M. Filter.
Sig.—Sixty minims contain seven and a half grains.
Bartholow.

℞ Quiniæ bisulphat.....gr. 50
Acid. sulphur. dil.....♏ 100
Aquæ font..... ℥ j.
Acid. carbol. liq.....♏ v.
Solve.
The quinine is dissolved by the aid of heat and after filtration the carbolic acid is added.

Lente's Solution of Quinia.

℞ Quiniæ sulphat.....gr. 80
Aquæ.....fl. ℥ 1
Acid. sulphur. dil.....q. s.
Heat to boiling and add:
Acidi carbolici.....gr. 5
For hypodermic use. *Dr. F. D. Lente*

Cystitis.

℞ Acidi benzoici.....
Sodii biboratis.....aa gr. x.
Inf. buchu..... ℥ ij.

This amount three or four times a day.

Dr. A. J. C. Skene.

This may almost be called specific in its influence in the earlier stages of cystitis, affording rapid and lasting relief. The diet should be carefully regulated, and the skin and bowels kept in active condition.

As an Aphrodisiac.

℞ Ergotin (aq. ext.).....ʒ j.
Sanguinariæ.....gr. ij.

Ft. pil. No. xx. M. Sig.—One three times a day.

Or the following:

℞ Tinct. sanguinariæ..... ʒ iij.
Ext. stillingiæ fl..... ʒ v.

M. Sig.—Fifteen to twenty drops in water three times a day. *Dr. Bartholow.*

Impotence with Spermatorrhæa.

℞ Tinct. canthar.....gtt. vj.
Tinct. ferri chlor.....gtt. xv–xx.

Sig.—Thrice daily in water. *Dr. H. C. Wood.*

This prescription has been found to cure the above condition so speedily as to commend itself to the use of all medical men in the treatment of these cases.

Treatment of Seminal Emissions.

Bumstead gives the following prescription for its special tonic effect upon the genital organs:

℞ Tr. ferri chloridi..... ..... ℥ iii.
Ext. ergot. fld. (Squibb's)..... ℥ iii.
M. Et. sig.—A teaspoonful in water after each meal.

As a direct means of diminishing the frequency of the emissions, B. recommends:

℞ Potass. bromidi..... ℥ i.
Tr. ferri chloridi..... ℥ i.
Aquæ..... ℥ iii.
M. Et. sig.—From one to two teaspoonfuls in water, after each meal, and at bedtime.

The avoidance of tobacco in all its forms, cleanliness of mind and body, laxatives when needed, and, in a word, attention to the rules of hygiene, are to be strictly enjoined.

Elixir Chloroformi Compositus.

Prof. W. F. McNutt has been in the habit for several years of prescribing Collis Browne's chlorodyne in certain cases of asthma, colic, diarrhœa, neuralgia, rheumatism, hysteria, etc. He states that it has seldom failed to be of some benefit, and has often acted like a charm; in fact, it is a most excellent and reliable anodyne, anti-spasmodic and sedative.

On account of several objections to its use, I have,

after a great deal of experimentation, adopted the following formula as a substitute for chlorodyne, viz:

℞ Morph. mur.....gr. ½
Chloral hyd.....
Chloroform.....aa ʒ ss.
Tinct. cannab. ind.....
Tinct. capsici.....
Acid. hydrocyan. dil.....aa m. xx.
Spt. menth. pip.....♏ x.
Syr. sassafras co. ad..... ℥ j.

Dose: ʒ j.

This I have named Elixir Chloroformi Compositus, and can heartily recommend it to those who have been in the habit of using chlorodyne. To those who have never used chlorodyne, I may say that they will find elix. chlorof. comp. a most efficient remedy for many purposes and under many circumstances; for instance, in whooping-cough, asthma, emphysema, cough of many phthisical patients, in many cases of hysteria, and especially in many cases of dysmenorrhœa, it certainly has no equal. Given as an anodyne, it seldom produces headache or disturbance of the digestion, as does morphine; or depresses the heart's action, as does hydrate of chloral. In diarrhœa accompanied with tormina, in teaspoonful doses repeated every two or three hours, it generally acts quickly and satisfactorily.

In many cases of diarrhœa in children, a few drops of the elixir, together with a few drops of castor oil and vini ipecac, in syrup of acacia, make a most efficient remedy. *Prof. W. F. McNutt.*

Elixir Simplex.

℞ Spiritus aurantii (1 in 10).....fl. ʒ 2
" cinnamomi (1 in 10).....♏ 10
Alcoholis.....fl, ℥ 6
Syrupi.....fl. ℥ 6
Aquæ.....fl. ℥ 6

This Elixir may be made the vehicle of various remedies which have an unpleasant taste, or are otherwise not readily taken. This form of administration, however, should be used *very* sparingly and judiciously to prevent patients acquiring a taste for "cordials" and alcoholic beverages, generally.

Pulvis Morphiæ Co.—(Tully's Powder.)

℞ Morphiæ sulph.....gr. 1
Camphoræ.....
Pulv. glycyrrhizæ.....
Calcii carbonat.....aa gr. 20

Mix. 10 grains contain $\frac{1}{6}$ grain of morphia sulphate. This formula is taken from Dr. Tully's Materia Medica. (Springfield, 1858, vol. 1, page 153.)

Chronic Alcoholism.

℞ Zinci oxidi..... ʒ j.
Piperin..... ℈ j.
Ft. pil. No. xx.

M. Sig.—One pill three or four times a day.

Dr. Bartholow has found that this will relieve the gastric catarrh and trembling, and diminish the craving for strong drink in this condition.

Emetic in Narcotic Poisoning.

℞ Cupri sulphat.....gr. vj.
Aquæ destillat..... ℥ ij.

M. Sig.—A tablespoonful every fifteen minutes till vomiting ensues. *Bartholow.*

In Internal Piles.

℞ Tinct. ferri perchlor.....♏ x.
Aquæ..... ℥ j.

Fiat injectio. *Mr. Erichsen.*

In Fissure or Ulcer of Anus.

℞ Ext. belladonnæ.....
Plumbi acet.....āā gr. ij.
Tannin.....gr. iv.
Sevi præp.....q. s.

Fiat suppositorium. *Mr. Erichsen.*

In Indurated Glands Following Scarlet Fever.

℞ Calcii sulphid.....gr. j.
Sacchar. lactis.....gr. x.

Misce. Fiat pul. x. One powder every hour or two. *Dr. S. Ringer.*

In Eczema.

℞ Syr. tolu.....
Vini ferri.....āā ℥ ss.
Liq. arsenicalis.....♏ xij.
Aquæ anethi..... ℥ j.

Misce. A teaspoonful thrice daily, directly after food—for a child two years old.

Sir Erasmus Wilson.

In Scrofulous and Tuberculous Glands.

℞ Calcii sulphid.....gr. j.
Aquæ..... ℥ x.
Misce. A teaspoonful every hour.

Dr. S. Ringer.

In Chronic Eczema of the Aged.

℞ Inf. cinchonæ.....f ℥ vj.
Liq. calcis.....f ℥ ixss.
Tinct. lupuli.....
Succi conii.....ana f ʒ ij.
Misce. A wineglassful three times a day.

Dr. Neligan.

In Irritable Bladder, with Alkaline Urine.

℞ Acidi nitrici dil.....f ʒ j.
Acidi hydrochlor. dil.....f ʒ ss.
Aquæ destil.....f ℥ viij.
Misce. Sumat cochl. ij. ter die. *Mr. Coulson.*

In Acne with Plethora.

℞ Magnes. sulph..... ʒ ss.
Acid. sulph. aromat..... ♏ xx.
Ferri sulph.....gr. iij.
Quiniæ sulph.....gr. j.
Vini colchici..... ♏ x.
Syr. zingiberis..... ʒ j.
Aquæ..... ℥ j.
Fiat haustus. To be taken twice or thrice a day, with an aperient pill if needed. *Dr. Tilbury Fox.*

In Chronic Skin Diseases, Etc.

℞ Sodæ arseniatis.....gr. ij.
Aquæ destil. q. s. Solve, et adde:
Guaiaci pulv..... ʒ ss.
Antim. sulphurati..... ℈ j.
Mucil. acaciæ.....q. s
Misce caute et div. in pil. xxiv. *Sir E. Wilson.*

In Retention of Urine.

℞ Magnes. sulph.....gr. xxx.
Potass. bicarb.....gr. xx.
Potass. nitratis.....gr. x.
Aquæ..... ℥ j.
Misce. Fiat haustus. *Sir James Paget.*

In Simple Erysipelas.

℞ Tinct. ferri perchlor..... ♏ xx.
Aquæ..... ℥ j.
Fiat haustus. To be given every third or fourth hour. *Mr. Erichsen.*

In Hemorrhages.

℞ Ergotin.....gr. iij.
Sp. rectificat.....
Glycerini.....ana ♏ ij.
Fiat injectio. For subcutaneous injection.
Langenbeck.

In Oxaluria, with Debility and Irritability.

℞ Acidi nitrici dil.....
Acidi hydrochlor. dil.....ana ♏ v.
Inf. serpentariæ..... ℥ j.
Fiat haustus ter die sumendus. *Dr. G. Bird.*

Anaphrodisiac.

℞ Camphoræ.....
Ext. latucæ.....ana ℈ iiss.
Misce. Fiant pil. xx. From four to six pills to be taken daily. *M. Ricord.*

Ichthyosis.

℞ Adipis benz..... 1 ounce.
Glycerinæ.....40 minims.
Ung. petrolei..... ..... ½ ounce.
M.—Sig. Apply daily after washing or bathing.
Dr. Louis A. Duhring, Phila.

Eczema Infantile.—Ointment.

℞ Salicilic acid..... 2 parts.
Subnitrate of bismuth..... 30 "
Corn starch..... 20 "
M. Ointment of rose-water.....100 "

Spread thickly on stout pieces of muslin and cover carefully every portion of inflamed and moist skin. This should not be used on the scalp, in consequence of matting the hair. For the scalp use oily applications.

℞ Oil of cade, and Almond oil equal parts. If thick crust is present, remove by poulticing—wear an oiled silk skull cap.

Dr. George Henry Fox, New York.

Diabetes Mellitus.

℞ Sodii salicylat.....ʒiij.
Liq. potass. arsenitis.....ʒj.
Glycerinæ.....℥j.
Aq. cinnam.....ad ℥iij.

M. Sig.—Dessertspoonful three times a day.

J. C. Wilson.

Sea Sickness.

℞ Chloroform.....℥ss.

M. Sig.—Two to five minims on sugar every half hour until relieved. *Bartholow.*

Conjunctivitis.

℞ Zinci sulphat.....gr. ss.
Sodii biborat.....gr. ij.
Aq. camphoræ.....
Aquæ.....aa ʒij.

Filter.

M. Sig.—Two or three drops in the eyes twice or three times daily. *Dixon.*

For Inflammation.

℞ Ichthyol.....gr. xlv.
Plumbi iodidi.....gr. xlv.
Ammonii chloridi.....gr. x.
Petrolati.....ad ℥i.

M. Sig.—Rub gently into the site of the inflammation every hour. This is effective in relieving deep-seated as well as superficial inflammation.

Dr. W. T. Slevin.

Hemorrhage.

℞ Ext. suprarenal gland..... ʒi.
Sol. resorcin (one per cent.)..... ℥j.
Mix and filter.

Sig —Apply by swab or spray to the site of the hemorrhage. *Dr. S. Oppenheimer.*

Suprarenal extract is unquestionably the most potent means of arresting *epistaxis* and other forms of hemorrhage at our command. In emergency the solution may be made by stirring up five to ten grains of the powder in a drachm of water. This solution, however, deteriorates very rapidly, but the prescription here given makes a solution that keeps for several weeks. The small amount of resorcin does not diminish the activity of the suprarenal and is not irritating.

The above solution is one of the most effective measures for treating *Hay Fever.* It is to be used as a spray several times a day, with or without the addition of cocaine.

Pyæmia.

℞ Acid. salicylici..... ℥ss.
Sodii biborat..... ʒj.
Glycerinæ..... ℥j.
Aq. menthæ pip..... ℥v.
M. Sig —Tablespoonful every two or three hours.
Bartholow.

Varicose Veins.

℞ Ex. hamamelis fl..... ℥ij.
M. Sig.—Teaspoonful three or four times a day, with compresses applied externally. *J. V. Shoemaker.*

Cystitis.

℞ Tr. hyoscyomi.....♏. xx.
Potassii bicarb.....gr. v.
Infus. buchu .....ad ℥ ij.

M. Sig.—This amount after meals and at bedtime when the urine is *acid.* The equivalent of this prescription is now made into tablet form by several pharmaceutical chemists.

℞ Ac. benzoici.....gr. iij.
Sodii biborat.....gr. ij.
Ext. buchu.....gr. j.
Ext. tritici repentis.....gr. j.
Ext. hydrangiæ.....gr. ss.
Atropiæ sulphat.....gr. $\frac{1}{500}$

M. Ft. tablet.

Sig.—One or two after meals and at bedtime when the urine is *alkaline.*

Acute Cystitis.

℞ Tr. aconit.....℥ j.
Spt. æther. nitros.....℥ j.
Liq. potass. citrat.....q. s. ad ℥ vj.

M. Sig.—Dessertspoonful every four hours until all fever ceases and the pulse is quiet. *Hare.*

Head Lice.

℞ Sodii hyposulphitis.....℥ ij.
Acid. sulphurosi dil.....℥ iv.
Aquæ.....q. s. ad ℥ xvi.

M. Sig.—Apply once daily. *Startin.*

Toothache.

℞ Creasoti.....ʒij.

M. Sig.—Moisten a very small pledget of cotton and lay it in the carious cavity; then pack a larger piece of plain cotton over it to retain it. *Henson.*

OR

℞ Acid. arseniosi.....
Cocaini muriat.....aa gr. xv.
Menthol cryst.....gr. iiss.
Glycerinæ.....f ʒiij.

M. Sig.—A pledget of cotton moistened with this, and placed in the cavity of the tooth, will quickly check the pain. *L' Union Médicale.*

Painless Tooth Extraction.

℞ Cocaini hydrochlor.....
Morphiæ sulphat.....
Chloral hydrat.....
Acid. carbolic.....aa gr. x.
Aq. rosæ.....ʒx.

M. Sig.—Inject with a hypodermic syringe into the gums.

Diabetes Insipidus.

℞ Sodii salicylat.....ʒiv.
Glycerinæ.....℥ij.
Aquæ.....q. s. ad ℥iij.

M. Sig.—Two teaspoonfuls three times daily.

Da Costa.

Alcoholism.

℞ Tr. nucis vomicæ.... .....♏lxxx.
Tr. gentian co..... ....
Tr. columbæ co.....aa ℥ij.

M. Sig.—Dessertspoonful before each meal, in water.
Loomis.

OR

℞ Tr. capsici..... ....ʒiv.
Potass. bromid..... ...ʒiv.
Spt. ammon. aromat.....℥iiss.
Aq. camphoræ.....ad ℥vj.

M. Sig.—Dessertspoonful several times daily for the depression. *Ringer.*

Hiccough.

℞ Apomorphiæ muriat.....gr. $\frac{1}{10}$
Aquæ.....♏x.

M. Sig.—Inject hypodermically.....*Ringer.*

Tapeworm.

℞ Ol. filicis maris æther.....ʒii–iij.
Emuls. amygdal. dulc.....ad ℥vj.

M. Sig.—In the evening a light meal is eaten. At bedtime, about twenty minutes apart from each other, this medicine is taken in two doses. The next morning early, about five o clock, two tablespoonfuls of castor oil are administered, and these followed about an hour later by another tablespoonful. *Hugo Engel.*

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

FAVORITE PERSCRIPTIONS.

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

For By Dr.

℞

Dose

How often

Remarks

INDEX.

Abdomen, painful affections of, 116.
Acetanilid, in relief of pain, 102.
Acne, 195, 197.
    indurata, 195.
    intermenstrual, 200.
    obstinate, 198.
    rosacea, 202.
    with plethora, 218.
Adenitis of children, 138.
Alcoholism, 225.
    chronic, 216.
Alkaline bath for pruritus, 203.
    laxative (rhubarb and soda), 50, 51.
    mixture for nursing children, 125.
    urine, 218.
Aloes mixture, compound, 47.
Amenorrhoea, 119.
    from anaemia, 190.
    " chlorosis, 190.
    " uterine atony, 115.
    with anaemia, 110.
Ammonia mixture, carbonate, 22.
    mixture, chloride, 23.
Amygdalitis, 34.
Anaemia, 182.
    (Blaud's pills), 177, 182.
    amenorrhoea from, 190.
    in malaria, 174.
    of chlorosis, 114.
    of malaria, 154.
    with amenorrhoea, 110.
Anaemic headache, 100.
Anaesthetic application, 199.
    in labor, 117.
Anaphrodisiac, 220.
Angina pectoris, 31.
    (heart stimulant), 37.
Anodyne (elix. chlorof. co.), 214.
    Hoffman's, and iodide of potash, 25, 28.
Anorexia, 58.
Anthelmintic, 62, 107, 143, 225.
Anti-emetic mixture, 49.
Anti-epileptic mixture, 74.
    " " pill, 75.
Antifermentative, 66.
Antipyrin in headaches, 102.
Antirheumatic, 157, 158.
    in pericarditis, 31.
    pills, 150.
Antitoxin in croup, 148.
    treatment of diphtheria, 146, 147, 148.
    immunizing doses of, 148.
Anus, fissure of, 217.
    pruritus of, 197.
    ulcer of, 217.
Aperient, ferruginous, 175.
    for children, 137.
    in gout, 161.
    pills, 54.
Aphrodisiac, 213.
Appetite, loss of, 58.
Apthae, 36, 142.
    of phthisis, 14.
Ascaris lumbricoides, 127.
    vermicularis, 127.
Asthenic inflammatory fever, 160.
Asthma in children, 144.
    hay, 12, 13.
    spasmodic, 11, 13.
Asthmatic fits of emphysema, 9.
    mixture, 22.
    paroxysm, 10.
Atonic dyspepsia, 62.
Axilla, odors from, 192.

Barker's post partum pills, 114.
Basham's mixture, 69, 176.
Bedsores, 197, 204.
Bell's gargle, 17.

Belladonna mixture for epilepsy, 77.
Bicarbonate of soda, 50.
Bismuth and Dover's Powders, 56
Bismuth and pepsin powder, 56, 57.
Bitters, Smith's, 175.
Bladder, irritable, 120, 218.
Blaud's Pills, 177, 182.
Bleeding, 15, 189, 219, 222.
passive, 132.
suprarenal extract for, 222.
hemorrhoids, 188.
Breast, painful affections of, 116.
Breath, foul, 192.
Bronchial catarrh, 10.
acute, 34, 36.
Bronchitis, 19, 22, 23, 30, 37, 38.
acute, 14.
" catarrhal, 29.
" in children, 123.
catarrhal of infants, 125.
in children, 128–132.
Bronchocele, 34.
Brown mixture, 23.
Brown-Sequard's Neuralgic Pills, 79.
Bruises (contusions), 203.
Burns, 194.
glue mixture for, 187.

Cachexia of chronic rheumatism, 159.
malarial, tonic in, 173.
of rheumatic gout, 159.
syphilitic, 166, 169.
Cancrum oris, 32.
Cannabis mixture, 76.
Carbonate of ammonia mixture, 22,23.
Cardiac debility, 31, 32, 33, 34.
" with scanty urine, 17.
dilatation, 16, 18.
disease, dyspnoea of, 29.
dropsy, diuretic in, 9.
weakness and dilatation, 16.
Carron oil, 194.
Carson's paint, 192.
Caseous pneumonia, 20.
Castor oil mixture, 49.
Catarrh, acute, 11, 19.
" bronchial, 34, 36.
bronchial, 10.
Catarrh, chronic bronchial, 37.
" gastric, 59, 65.
epidemic (influenza), 17, 18.
gastric, 44, 64.
summer (hay asthma), 12, 13.
Catarrhal bronchitis, acute, 29.
of children, 125.
Cathartic pills, vegetable, 55.
Cerebral congestion, 74.
tumors, headache from, 101.
Cervical endometritis, chronic, 109.
Cervicitis, chronic, 110.
Cervix uteri, ulceration of, 117.
Change of life, laxative in, 118.
Chilblains, 185, 192, 196, 203.
Children, acute bronchitis in, 123.
adenitis of, 138.
alkaline mixture for, 125.
asthma in, 144.
bronchitis in, 128—132.
chronic diarrhoea of, 133.
convulsions in, 143.
delirium of, 127.
diarrhoeal mixture for, 123, 127.
earache in, 144.
eczema of, 140, 220.
expectorant mixture for, 20.
febrile coughs of, 137.
fever in, 143.
irritable stomach of, 135.
itch in, 133, 134.
laxative alkaline, for, 143.
nocturnal incontinence of, 124, 141.
pleurisy in, 142.
porrigo in, 136.
scabies in, 133, 134.
severe coughs of, 136.
summer diarrhoeas of, 49, 124, 144, 145.
tapeworm in, 143.
tonic for, 139.
vomiting of, 135.
Chloasma of pregnancy, 120.
(tinea versicolor), 201.
Chlorate of potash mixture, 26.
Chloride of ammonium mixture, 23.
Chlorosis, 115.
(Blaud's pills), 177, 182.

Chlorosis, amenorrhoea from, 190.
anaemia of, 114.
Cholera infantum, 126.
mixture, Squibb's, 53.
morbus, 45.
Chordee, 168.
Chorea, 72, 76, 136, 141.
Chrysophanic acid ointment, 194.
Cirrhosis, first stages of, 47.
Climacteric, menorrhagia of, 114.
metrorrhagia of, 115.
Cod liver oil emulsion, 207, 209.
phosphorated, 207.
phosphorated, 208.
for children, 209.
emulsion of, with lime, 209.
mixture, 209.
Cold, acute, 20, 37.
Colic, 64, 67.
flatulent, 61.
lead, 43.
Compound aloes mixture, 47.
licorice mixture, 23.
licorice powder, 56.
rhubarb mixture, 50.
syrup of hypophosphites, 77.
Congestion of larynx, 36.
acute, of larynx, 33.
" of pharynx, 33.
cerebral, 74.
laryngeal, stimulant in, 36.
Congestive dysmenorrhoea, 109.
headache, 99.
Conium mixture for epilepsy, 78.
Conjunctivitis, 192, 221.
Constipation, 46, 63.
aloes and fellis pills for, 54.
habitual, 45.
(see laxatives.)
Consumptive cough, 32.
Continued fever, 149.
Contusions, 203.
Convulsions, infantile, 81, 126, 143.
Coryza, 20, 37, 40, 41.
acute and hay asthma, 12.
syphilitic, 170.
Cough, 21, 38.
consumptive, 32, 38.
Cough, febrile of children, 137.
of infants, 125.
of elderly people, 38.
severe of children, 136.
spasmodic, 32.
due to bronchitis, 38.
" " laryngitis, 38.
" " phthisis, 38.
whooping, 15, 132, 133, 136, 139.
Counter irritant in pleurisy, 28.
Cretinism, 138.
Croton oil liniment, 193.
Croup, diphtheritic, 9.
" 148.
stimulant in, 136.
Cyanide and bromide of potash, 27.
Cystitis, 213, 223.
acute, 223.

Debility, cardiac, 31, 32, 33, 34.
cardiac, with scanty urine, 17.
and irritability in oxaluria, 220.
and loss of appetite, 58.
Delirium, 75.
of children, 127.
of fever, 75
of typhoid, 149.
of typhus fever, 158.
Depilatory, 195.
Diabetes, insipidus, 224.
mellitus, 221.
Diarrhoea, 46, 47, 48, 58, 135.
acute, 68.
bismuth, chalk and opium for, 53.
bismuth and Dover's powder for, 56.
castor oil mixture for, 49.
children's, 123, 127.
chronic, 46, 132.
of children, 133.
and dysentery, 59, 64.
hot drops for, 44.
opium, rhubarb and camphor, 44.
of phthisis, 53.
powder, 57.
rhubarb and lime for, 51.
simple, of infants, 135.
summer, of children, 124, 144, 145.

Diarrhoea of typhoid, 153, 162.
West's mixture for, 44.
Dilatation of heart, 16, 37.
Dinner pill, 60.
Dipsomania, 80.
Diphtheria, 146, 147, 148.
antitoxin treatment of, 147.
Letzerich's treatment, 15.
Diphtheritic croup, 9.
" " antitoxin in, 148.
Diuretic, 48, 68.
in cardiac dropsy, 9.
in pleurisy, 29.
pill in pleurisy, 29.
Dressing for wounds, 204.
Drink, craving for, 80.
Dropsy, cardiac, diuretic in, 9.
of emphysema, 36.
pleural, 28.
Dysentery, 59, 64, 65, 68.
Hope's mixture for, 50.
acute, 45.
" and chronic, 51.
Dysmenorrhoea, 117, 118.
congestive, 109.
obstinate, 116.
rheumatic, 108, 112.
Dyspepsia, atonic, 62.
(bicarbonate of soda), 50.
(bismuth and pepsin), 56, 57.
(compound bismuth powders), 56.
(compound rhubarb mixture), 50.
fermentative, 65.
flatulent, 61.
irritative and gastralgia, 51, 62.
(rhubarb and lime), 51.
(rhubarb and soda) 50, 51.
Dyspeptic headache, 98.
Dyspnoea of cardiac disease, 29.
emetic in laryngeal, 35.

Ear, furunculosis of, 41.
Earache, 41, 144.
Eczema, 193, 199, 201, 217.
acute, 195.
chronic, of the aged, 218.
infantile, 220.
of children, 140, 217, 220.
Eczema of hands, 201.
Elix. chlorof. comp. (McNutt), 215.
simplex, 216.
Emetic in dyspnoea of laryngitis, 35.
in narcotic poisoning, 217.
Emissions, seminal, 214.
Emphysema, 10, 13, 29.
asthmatic fits of, 9.
dropsy of, 36.
Emulsion, cod liver oil, 207—209.
expectorant, 24.
Endometritis, chronic cervical, 109.
Enuresis, 124, 141.
Epilepsy, idiopathic, 71, 74, 75, 80, 81, 105.
in the weak and anaemic, 7
(belladonna mixture), 77.
(conium mixture), 78.
(ergot mixture), 79.
Epistaxis, 39.
(hemorrhage), suprarenal extract for, 222.
Ergot mixture for epilepsy, 79.
Ergotin solution for hypodermic use, 211.
Eruptions, syphilitic, 194.
Erysipelas, 202, 219.
facial, 193.
Excoriations, 199.
Expectorant emulsion, 24.
for children, 20.
mixture, 19, 20, 21, 24, 26.
" Stoke's, 23.

Facial erysipelas, 193.
Febrile cough of children, 137.
Feet, fetid, 192, 204.
sweating, 204.
Ferruginous aperient, 175.
Fever, 150, 156.
acute, 154.
asthenic inflammatory, 160.
continued, 149.
delirium of, in children, 127.
" of typhus, 158.
hectic, 155.
in children, 143.
loss of hair after, 204.

Fever, milk, 111.
scarlet, 108, 117.
sthenic, 151.
symptomatic, 151.
typhoid, 159.
Fissured nipples, 185, 195, 198.
Flatulence, 62.
Flatulent colic, 61.
dyspepsia, 61.
(see dyspepsia.)
Foul breath, 192.
Freckles, lotion for, 196, 204.
Frostbite, 185, 192, 196, 203.
Furunculosis of the ear, 41.

Gargle, Bell's, 17.
Gastralgia, 49, 51.
Gastric catarrh, 44, 59, 64, 65
irritability, 61.
ulcer, 60.
chronic, 64.
Gastro-enteritis, 57.
Glands, indurated from scarlatina, 217.
scrofulous, 218.
tuberculous, 218.
Glue mixture for burns, 187.
Goa ointment, for skin affections, 188.
Goitre (bronchocele), 34.
Gonorrhoea, 167, 169.
acute stage, 168.
inflammatory stage, 167.
repeated attacks, 168.
second stage, 167.
Goodell's tonic formulas, 176.
Gout, 152, 156, 160, 161, 162.
aperient in, 161.
pills (White's), 159.
rheumatic in cachectic subjects, 159.
Gouty headache, 96.
Gravel (lithiasis), 152.
Griffith's mixture, 175.
Grip, coryza accompanying, 40.
Gross's neuralgic pills, 78.
Gums, ulcerated, 18.

Habitual constipation, 45.
Hall's mixture, modified, 176.
Haemoptysis, 19, 31, 67.
Hemostatic (Monsel's), 189.
Hair, loss of, after fever, 204.
Hammond's mixture, 75.
Hands, eczema of, 201.
Hay asthma, 12, 13.
Hay fever, suprarenal extract for, 222
Headache, 82, 102, 103.
(nervine tablets), 104.
anaemic, 100.
congestive, 99.
dyspeptic, 98.
from cerebral tumors, 101.
gouty, 96.
reflex, 74.
rheumatic, 96.
sick, 82.
uraemic, 96.
Head lice, 223.
Heart, dilated, 16, 37.
mitral regurgitation, 16.
disease, advanced, 32.
chronic, 16, 33.
enlarged without valvular lesion, 18.
disease, dyspnoea of, 29.
stimulant, 37.
Hebra's diachylon ointment, 193.
Hectic fever, 155.
Hemorrhage, 15, 189, 219, 222.
passive, 132.
suprarenal extract for, 222.
during pregnancy, 113.
Hemorrhoids, 69.
bleeding, 188.
external, 198.
internal, 217.
laxative for, 69.
Herpes, 199.
circinnatus, 197, 198.
Hiccough, 225.
Hoarseness, 30.
Hoffman's anodyne and iodide of potash, 25, 28.
Hope's mixture for dysentery, 50.
Hot drops, for diarrhoea, 44.
Hydrocyanic mixture, 26.
Hydrothorax, 28.

Hyperemia of kidney, 48.
Hypnotic, 72, 79.
Hypophosphites, compound syrup of, 77.
Hysteria, 81, 104.
(nervine tablets), 104.

Ichthyosis, 220.
Impetigo, 196, 197, 202.
Impotence, 213.
Incontinence, nocturnal in children, 124, 141.
Indigestion, 66.
painful, 66.
Indurated glands from scarlatina, 217.
Infants, catarrhal bronchitis of, 125.
cough mixture for, 125.
simple diarrhoea of, 135.
Infantile convulsions, 81, 126.
eczema, 220.
syphilis, 124.
Inflammation, 221.
of nipples, 191.
Influenza, 17.
Injuries of muscles, 160.
" tendons, 160.
Insomnia, 81, 105.
(nervine tablets), 104.
of typhus, 158.
Iodide of potassium, with Hoffman's anodyne, 25, 28.
Iritis, lotion for, 189.
Irritative dyspepsia, 51, 62.
Irritable bladder, 120, 218.
stomach of children, 135.
Itch, 196, 197.
in children, 133, 134.

Kidneys, acute nephritis, 63, 65.
hyperaemia of the, 48.
inflammation of, 63.

Labor, anaesthetic in, 117.
Lactophenin in headache, 103.
Laryngismus stridulus, 133.
Laryngitis, acute, 33.
emetic in dyspnoea of, 35.
cough, due to, 38.
Laxative, alkaline, 60.
(rhubarb and soda), 50, 51.
mixture for children, 143.
compound licorice powder, 56.
for hemorrhoids, 69.
in change of life, 118.
pills, 54.
pills, Cole's, 53.
pills, Squibb's, 55.
in uterine disorders, 112.
Larynx, acute congestion of, 33.
stimulant in congestion of, 36.
ulceration of, 30.
weakness of, 30.
Lead colic, 43.
Letzerich's treatment of diphtheria, 15.
Leucorrhea and chronic cervicitis, 110.
Lice, head, 223.
Licorice mixture, 25.
mixture, compound, 23.
Liquid pepsin, 55.
Liquor chloroform co., 211.
Lithiasis, 152.
Liver, cirrhosis of, 47.
sluggish, 52.
Locomotor ataxia, 103.
acetanilid in, 102.
Loomis's tonic, 180.
Loss of hair after fever, 204.
Lotion, Taylor's, 191.

Malaria, acetanilid in neuralgia of, 102.
anaemia in, 174.
chronic, 151.
enlargement of spleen, 153.
anaemia of, 154.
pulvis (pulvis quiniae comp.) 158.
tonic in, 173, 183.
tonic for anaemia in, 174.
Mania, sub-acute, 73.
acute, 104.
Menopause, laxative at, 118.
Menorrhagia, 118.
of climacteric, 114.
Mercurial salivation, 14, 18.
Metritis, 120.
digestive tonic in, 174.

Metrorrhagia, 113, 115.
of climacteric, 115.
Migraine, 82, 102.
milk fever, 111.
Mistura ferri comp., 175.
Mitral regurgitation, 16.
Monomania of suspicion, 73.
Mouth, apthae of, 36, 142.
ulcerated, 18.
Muscular rheumatism, 155.
pains, chronic, 193.
Muscles injuries of, 160.

Narcotic, 79.
poisoning, emetic in, 217.
Nausea of uterine irritation, 116.
(see vomiting), 67.
(anti-emetic mixture), 49.
Nephritis, chronic interstitial, 52.
second stage, 63.
acute, 63, 65.
Nervine tonic, 75.
tablets, 104.
Neuralgia, 71, 72, 78, 102, 103.
tonic in, 80.
antipyrine in, 102.
acetanilid in, 102.
ovarian (nervine tablets), 104.
Neuralgic pills, Gross's, 78.
pills, Brown-Sequard's, 79.
Neuralgic rheumatism, 160.
Neurasthenia, 120.
(nervine tablets), 104.
Neuritis, acetanilid in, 102.
Night cough, 38.
Nipples, fissured, 185.
sore, 189, 195, 198.
inflammation of, 191.
Nitrate of potash mixture, 25e.
Nocturnal incontinence in children, 124, 141.

Odors from axilla, 192.
Opium, rhubarb and camphor in diarrhoea, 44.
Otitis, 41.
Otorrhoea, 41.
Oxaluria, 220.
Oxyuris vermicularis, 127.
Ozaena, 34, 35.

Painful affections of abdomen, 116.
" " breast, 116.
Painless tooth extraction, 224.
Parasites (head lice), 223.
Parasitic skin diseases, 185, 187.
Pediculosis, 146, 148, 223.
(ung. hydrarg. oxidi), 186, 187.
(head lice), 223.
(parasitic skin diseases), 187.
Pepsin, liquid, 55.
Pericarditis, anti-rheumatic in, 31.
Pertussis, 15, 132, 133, 136, 139.
Phagedena, sloughing, 197.
Pharyngitis, dry, 39
Pharynx, acute congestion of, 33.
relaxation of, 35.
Phenacetin, in relief of pain, 103.
Phosphorated cod liver oil, 208.
" " emulsion, 207.
Phosphorus, tincture of, 76.
mixture, 77.
Phthisis, 13.
apthae of, 14.
chronic, 11.
cough of, 38.
diarrhoea of, 53.
for expectorant and to destroy odor, 40.
rheumatism of, 156.
Piles, 69.
bleeding, 188.
external, 198.
internal, 217,
laxative for, 69.
Pitting of vaccinia, 200.
Plaster (McCready's capsicum), 189.
Pleural dropsy, 28.
Pleurisy, 35.
acute, 18, 30.
chronic, 33.
counter-irritant in, 28, 192.
" (Carson's paint), 192.
diuretic in, 29.
" pill in, 29.
in children, 142.

Pleurisy, sub acute, 8.
vesicant in, 192.
Pneumonia, 7, 27.
caseous, 20.
Poisoning, emetic in narcotic, 217.
Porrigo, 136.
Post-partum pills, Barker's, 114.
Potash chlorate mixture, 26.
cyanide and bromide mixture, 27.
nitrate mixture, 25.
Powder, Tully's, 216.
Pregnancy, chloasma of, 120.
hemorrhage during, 113.
pruritus of, 188.
sick stomach of, 110.
vomiting of, 109, 110, 111, 119.
Prurigo senilis, 198.
Pruritus, 186, 200, 202.
alkaline bath for, 203.
ani, 197.
of pregnancy, 188.
vulvae, 117, 190.
Purgative, 59.
Purge, strong, 63.
Putrid sore throat of scarlatina, 35.
Pyaemia, 222.

Quinine for hypodermic use, 212.
Lente's solution of, 212.
Quinsy, 39.

Rectum, ulcers of, 199.
Reflex headache, 74.
Regurgitation, mitral, 16.
Relaxation of pharynx, 35.
Renal hyperaemia, 48.
Retention of urine, 219.
Rheumatic dysmenorrhoea, 108, 112.
gout in cachectic subjects, 159.
headache, 96.
pain, application for, 150.
Rheumatism, 150, 155, 157, 158, 161.
acute, 154.
chronic, 152, 162.
" of cachectic subjects, 159.
liniment for, 162.
muscular, 155.
neuralgic, 160.
Rheumatism of phthisis, 156.
(salicylic acid), 157, 161.
Rhinitis, 40, 41.
Rhubarb and lime, 51.
and soda, 50, 51.
Rickets, 142.
Ringworm, 197, 198.
Round worm, 127.
Rum stomach, 43.

Salicylic acid in rheumatism, 157, 161.
Salivation, mercurial, 14, 18.
Sarcina ventriculi, 62.
Scabies, 196, 197.
in children, 133.
Scarlatina, 128, 137.
indurated glands following, 217.
putrid sore throat of 35.
Scrofulous glands, 218.
Sea sickness, 221.
Sedative mixture, 24, 67, 78.
in uterine affections, 118.
Seminal emissions, 214.
Sick headache, 82.
Simplex elixir, 216.
Skin diseases, chronic, 219.
parasitic, 185, 187.
(ung. hydrarg. ox. rub.), 186, 187.
(sulphur paste), 186.
(ung. picis alk.), 187.
(chrysophanic acid ointment), 194.
(Goa ointment), 188.
Sleeplessness in uterine disorders. 107.
(insomnia), 81.
(hypnotic), 72, 79.
Sloughing phagedena, 197.
Smith's bitters, 175.
Soda bicarbonate mixture, 50.
Sore throat, 39.
putrid, 35.
syphilitic, 171.
(quinsy), 39.
Sore nipples, 189, 195, 198.
Simple elixir, 216.
Spasmodic asthma, 11, 13.
cough, 32.
Spermatorrhea, 213.
Spleen, malarial enlargement of, 153.

Squibb's cholera mixture, 53.
laxative pills, 55.
Sthenic fever, 151.
Stimulant in croup, 136.
in congestion of larynx, 36.
Stimulating expectorant, 19, 26.
Stoke's expectorant, 23.
Stomach (acute gastric catarrh), 64.
(chronic gastric catarrh), 59, 65.
(compound rhubarb mixture), 50.
(gastralgia), 49, 51.
(gastric irritability), 61.
(gastric ulcer), 60, 64.
irritable, of children, 135.
rum, 43
(sarcina ventriculi), 62.
Stomatitis, 137.
Strangury, 170.
Strychnia mixture, 176.
Strychnine tonic. 181.
St. Vitus's dance, 72, 76, 136, 141.
Sulphur paste, 186.
Sunburn, lotion for, 197, 204.
Suprarenal extract for hemorrhage, and hay fever, 222.
Suspicion, monomania of, 73.
Sympotomatic fevers, 151.
Syphilis, 169, 170, 171.
(bichloride of mercury), 166.
(iodide of ammonium), 168.
(iodide of potassium), 166.
mixed treatment, 170, 171.
(syrup mercury biniodide), 165.
(Taylor's mixed treatment), 165.
(Thompson's mixed treatment), 165.
early stage, 168.
infantile, 124.
rebellious cases, 171.
Syphilitic cachexia, 166, 169.
coryza, 170.
eruptions, lotion for, 194.
sore throat, 171.

Tan, lotion for, 196.
Tapeworm, 225.
in children, 143.
Taylor's lotion, 191.
Tendons, injuries of, 160.
Thomson's tonic, 181.
Thread worms, 127.
Throat affections, chronic, 12.
putrid sore, 35.
sore, 39.
ulcerated, 32.
Thrush, 36, 142.
Thyroid treatment of cretinism, 138.
Tincture of phosphorus, 76.
Tinea circinata (ringworm), 197, 198.
versicolor (chloasma), 120, 201.
Tobacco heart (heart stimulant), 37.
Tonic, digestive, 174.
and stimulant, 181.
for anaemia in malaria, 174.
for Bright's disease, Basham's, 69.
for children, 139.
formulas, 176, 177, 178.
(iodide mixture), 179.
(iron and cinchona), 179.
(iron and citrate of ammonia), 178.
(iron and quinine), 180.
(ferri, quinia et strychnia), 174.
(sulphate of cinchona), 179.
in malaria. 173, 183.
in neuralgia, 80.
lemon, 179.
(wine of iron and citrate of quinia), 178.
Loomis's, 180.
(elixir cinchona and iron), 180.
(wine of citrate of iron), 181.
muriatic acid mixture, 175.
(mistura ferri comp.), 175.
nervine, 75.
pills, 173.
preparations, 173.
Smith's bitters, 175.
(wine of iron and citrate of quinidiæ), 176.
strychnia mixture, 176.
strychnine, 181.
Thomson's, 181.
Tonsillitis, 34.
Toothache, 224.
Tooth extraction, painless, 224.

Tuberculous glands, 218.
Tully's powder, 216.
Tumors, cerebral, 101.
Typhoid, diarrhoea of, 153, 162.
delirium of, 149.
fever, 159.
Typhus, delirium of, 158.
insomnia of, 158.

Ulcer, gastric, 60.
chronic gastric, 64.
of anus, 217.
of rectum, 199.
Ulcerated gums, 18.
mouth, 18.
throat, 32.
Ulceration of the larynx, 30.
of cervix uteri, 117.
(sloughing phagedena) 197.
Uraemia, 52.
Uraemic headache, 96.
Urine, acid (cystitis), 223.
alkaline (cystitis), 223.
retention of, 219.
scanty, with cardiac debility, 16.
Urticaria, 202.
Uterine affections, sedative in, 118.
atony, amenorrhoea from, 115.
disorders, laxative in, 112.
" sleeplessness in, 107.
irritation, nausea of, 116.
tumor, 120.

Vaccinia, pitting of, 200.
Vaginismus, 119.
Vaginitis, 119.
Varicose veins, 222.
Vegetable cathartic pills, 55.
Veins, varicose, 222.
Vomiting, 67.
of children, 135,
of pregnancy, 109, 110, 111, 119.
Vulvae, pruritus of, 117, 190.
Weakness of larynx, 30.
West's mixture for diarrhoea, 44.
White's gout pills, 159.
Whooping cough, 15, 132, 133, 136, 139.
second stage, 132.
Women, painful affections of the breast or abdomen of, 116.
Worms, 62, 127, 143, 225.
Wounds, dressing for, 204.

June-15. 1901

No 25 - JUN 6 1901

LIBRARY OF CONGRESS
00026606330